

ESPEN practical and partially revised guideline : Clinical nutrition in the intensive care unit

Clinical Nutrition

Singer, Pierre; Blaser, Annika Reintam; Berger, Mette M.; Calder, Philip C.; Casaer, Michael et al

<https://doi.org/10.1016/j.clnu.2023.07.011>

This publication is made publicly available in the institutional repository of Wageningen University and Research, under the terms of article 25fa of the Dutch Copyright Act, also known as the Amendment Taverne.

Article 25fa states that the author of a short scientific work funded either wholly or partially by Dutch public funds is entitled to make that work publicly available for no consideration following a reasonable period of time after the work was first published, provided that clear reference is made to the source of the first publication of the work.

This publication is distributed using the principles as determined in the Association of Universities in the Netherlands (VSNU) 'Article 25fa implementation' project. According to these principles research outputs of researchers employed by Dutch Universities that comply with the legal requirements of Article 25fa of the Dutch Copyright Act are distributed online and free of cost or other barriers in institutional repositories. Research outputs are distributed six months after their first online publication in the original published version and with proper attribution to the source of the original publication.

You are permitted to download and use the publication for personal purposes. All rights remain with the author(s) and / or copyright owner(s) of this work. Any use of the publication or parts of it other than authorised under article 25fa of the Dutch Copyright act is prohibited. Wageningen University & Research and the author(s) of this publication shall not be held responsible or liable for any damages resulting from your (re)use of this publication.

For questions regarding the public availability of this publication please contact openaccess.library@wur.nl



ESPEN Guideline

ESPEN practical and partially revised guideline: Clinical nutrition in the intensive care unit



Pierre Singer^{a,*}, Annika Reintam Blaser^{b,c}, Mette M. Berger^d, Philip C. Calder^e,
Michael Casaer^f, Michael Hiesmayr^g, Konstantin Mayer^h,
Juan Carlos Montejo-Gonzalezⁱ, Claude Pichard^j, Jean-Charles Preiser^k,
Wojciech Szczeklik^l, Arthur R.H. van Zanten^m, Stephan C. Bischoffⁿ

^a Intensive Care Unit, Herzlia Medical Center and Department of General Intensive Care and Institute for Nutrition Research, Rabin Medical Center, Beilinson Hospital, Sackler School of Medicine, Tel Aviv University, Tel Aviv, and Intensive Care Unit, Herzlia Medical Center, Israel

^b Department of Anaesthesiology and Intensive Care, University of Tartu, Tartu, Estonia

^c Department of Intensive Care Medicine, Lucerne Cantonal Hospital, Lucerne, Switzerland

^d Faculty of Biology and Medicine, Lausanne University Hospital, Lausanne, Switzerland

^e School of Human Development and Health, Faculty of Medicine, University of Southampton and NIHR Southampton Biomedical Research Centre, University Hospital Southampton NHS Foundation Trust and University of Southampton, Southampton, United Kingdom

^f Clinical Department and Laboratory of Intensive Care Medicine, Catholic University Hospitals (UZLeuven) and Catholic University Leuven, Leuven, Belgium

^g Division Cardiac-, Thoracic-, Vascular Anaesthesia and Intensive Care, Medical University Vienna, Vienna, Austria

^h Department of Pneumology, Infectious Diseases and Sleep Medicine, St. Vincentius Kliniken gAG, Karlsruhe, Germany

ⁱ Instituto de Investigación Sanitaria "imas12" Hospital Universitario 12 de Octubre, Madrid, Spain

^j Department of Clinical Nutrition, Geneva University Hospital, Geneva, Switzerland

^k Medical Direction, Hôpital Universitaire de Bruxelles, Erasme Hospital, Université Libre de Bruxelles, Brussels, Belgium

^l Centre for Intensive Care and Perioperative Medicine, Jagiellonian University Medical College & Anesthesia and Intensive Care Department, 5th Military Hospital, Krakow, Poland

^m Department of Intensive Care, Gelderse Vallei Hospital, Ede, The Netherlands & Wageningen University & Research, Wageningen, the Netherlands

ⁿ Department of Nutritional Medicine/Prevention, University of Hohenheim, Stuttgart, Germany

ARTICLE INFO

Article history:

Received 8 June 2023

Accepted 10 July 2023

Keywords:

Intensive care

Nutrition

Enteral

Parenteral

Guidelines

Recommendations

ESPEN

SUMMARY

Following the new ESPEN Standard Operating Procedures, the previous 2019 guideline to provide best medical nutritional therapy to critically ill patients has been shortened and partially revised. Following this update, we propose this publication as a practical guideline based on the published scientific guideline, but shortened and illustrated by flow charts. The main goal of this practical guideline is to increase understanding and allow the practitioner to implement the Nutrition in the ICU guidelines. All the items discussed in the previous guidelines are included as well as special conditions.

© 2023 Elsevier Ltd and European Society for Clinical Nutrition and Metabolism. All rights reserved.

1. Introduction

The European Society for Clinical Nutrition and Metabolism (ESPEN) has published guidelines on nutrition in the intensive care unit (ICU) in 2006 (enteral nutrition (EN)) and 2009 (parenteral nutrition (PN)) [1,2]. Since then, the ESPEN

methodology has been upgraded to the “S3 guidelines level” described elsewhere [3] resulting in rigorous evidence-based and consensus-based recommendations that were published in 2019 [4]. The present guideline is a shortening and a partial revision of the previous guideline. The determination of the effect of nutrition alone on any possible outcome is complicated by the fact that the severity of illness and the number of comorbidities encountered among adult ICU patients is increasing [5]. Furthermore, the large heterogeneity of the ICU population potentially reduces the external validity of the recommendations,

* Corresponding author.

E-mail address: Pierre.singer@gmail.com (P. Singer).

Abbreviations

ASPEN	American Society for Parenteral and Enteral Nutrition
BIA	bioelectric impedance analysis
BMI	body mass index
DHA	docosahexaenoic acid
ECMO	extracorporeal membrane oxygenation
EE	energy expenditure
EFI	enteral feeding intolerance
EN	enteral nutrition
EPA	eicosapentaenoic acid
ESICM	European Society of Intensive Care Medicine
ESPEN	European Society for Clinical Nutrition and Metabolism
FA	fatty acid
GIDS	Gastrointestinal Dysfunction Score
GLIM	Global Leadership Initiative on Malnutrition

GLN	glutamine
GRV	gastric residual volume
HDL	high-density lipoprotein
ICU	intensive care unit
LCT	long chain triglyceride
MCT	medium chain triglyceride
MNA	mini-nutrition assessment
MUST	malnutrition universal screening tool
NRS	nutritional risk screening
NUTRIC	nutritional risk in critically ill
PN	parenteral nutrition
RCT	randomized controlled trial
REE	resting energy expenditure
SCCM	Society for Critical Care Medicine
SGA	subjective global assessment
SIGN	Scottish Intercollegiate Guidelines Network
SOP	standard operating procedure

which should be seen as a basis to support decisions made for each patient on an individual basis [6]. For now, a gap exists between nutritional practices and the previous guideline [7] and many available studies address only one or at most some of the specific aspects of nutritional therapy. In the current guideline, the timing, route, dose and composition of nutrition will be discussed and recommendations will be made recognizing that acute metabolic changes as well as prolonged energy and protein deficits play a major role in patient outcome. The goal is to achieve optimal nutritional support for ICU patients and to illuminate the gaps in knowledge in order to provide priorities for future clinical research.

2. Methodology

The present practical guideline consists of 56 recommendations and is based on the ESPEN guideline on clinical nutrition in the ICU [4]. The original guideline was shortened by focusing the commentaries on the evidence and literature on which the recommendations are based. Some of the recommendations themselves have been changed, and the presentation of the content has been transformed into graphical presentations. The original guideline was developed according to the standard operating procedure (SOP) for ESPEN guidelines and consensus papers [3].

This SOP is oriented on the methodology of the Scottish Intercollegiate Guidelines Network (SIGN). Literature was searched and graded 1–4 according to the evidence, and recommendations were created and graded into four classes (A/B/O/GPP).

All recommendations were agreed in a multistage consensus process, which resulted in a percentage of agreement (%). Six recommendations (former R21, R22, R33, R35, R36, R37), were revised and updated (R21 and R35) or merged (R36 and R37) into one within the preparation of this practical guideline according to the recent evidence. These recommendations were voted on within the working group, and the percentages of agreement for the respective recommendations are taken from this voting. One recommendation (former R26, now 30) also needs revision, but will be revised during the next revision process in the near future. The guideline process summary and dissemination were funded exclusively by ESPEN. For further details on methodology, see the full version of the ESPEN guideline [4] and the ESPEN SOP [3].

3. Recommendations (Fig. 1)

3.1. General recommendations (Fig. 2)

1) Statement 1: Every critically ill patient staying for more than 48 h in the ICU should be considered at risk for malnutrition. (S1, strong consensus, 96%)

No specific nutritional score for use in the ICU has been validated thus far. The existing nutritional screening tools (nutritional risk screening (NRS) 2002 [8] and the malnutrition universal screening tool (MUST) score [9]) have not been designed specifically for critically ill patients. NUTRIC (= nutritional risk in critically ill) is a novel risk assessment tool merely based on the severity of disease was proposed [10].

Rather than mortality, long-term functional tests might better reflect the benefit of a nutritional policy [11]. It appears that among all the screening tools, NRS 2002 and MUST have the strongest predictive value for mortality, and they are the easiest and quickest to calculate [12]. Validation of their utility for daily clinical practice and nutrition management has not been performed, and therefore only expert opinion can be expressed.

A pragmatic approach should be considered for patients at risk, such as those staying in the ICU for more than two days, requiring mechanical ventilation, with infection, who are underfed for more than five days, and/or presenting with a severe chronic disease.

2) Medical nutrition therapy shall be considered for all patients staying in the ICU, mainly for more than 48 h (Fig. 2). (R1, Grade GPP, strong consensus 100%)

Commentary

There are no studies directly addressing the effect of duration of starvation on outcome in critically ill patients. Such studies could be considered unethical as energy intake is a mainstay of survival. Since previous recommendations [1,2], a cut-off of 48 h for the initiation of early nutrition and contraindications to early EN has been better established [13]. Additionally, one study showed possible benefit of a further delay of PN if EN is not possible/tolerated [14]. A careful and progressive re-introduction of nutrition may limit the risk of refeeding syndrome, especially in patients who are severely malnourished or have been in a starved state before admission.




Fig. 1. ESPEN practical guideline: Clinical nutrition in the intensive care unit. Overview of the structure of the guideline. EN, enteral nutrition; ICU, intensive care unit; PN, parenteral nutrition.




Fig. 2. General recommendations. Recommendations are presented in green boxes. References to other panels in ellipse. Abbreviations: see Fig. 1.




Fig. 3. Early medical nutrition (enteral/parenteral) in the ICU patient. Abbreviations: see Fig. 1.

3) Oral diet shall be preferred over EN or PN in critically ill patients who are able to eat.
(R3, Grade GPP, strong consensus, 100%) (Fig. 2)

Commentary

For patients able to eat, this route should be preferred if the patient is able to cover 70% of his needs from day 3–7, without risks of vomiting or aspiration. This amount (i.e. 70% or more of the needs) is considered as adequate.

4) A general clinical assessment should be performed to assess malnutrition in the ICU, until a specific tool has been validated.

Remark:

General clinical assessment could include anamnesis, report of unintentional weight loss or decrease in physical performance before ICU admission, physical examination, general assessment of body composition, and muscle mass and strength, if possible.

(R2, Grade GPP, strong consensus 100%)

Commentary

Weight and body mass index (BMI) do not accurately reflect malnutrition. Loss of muscle and sarcopenia have to be detected in the ICU, including in obese patients. Frailty has been suggested as the burden of comorbid disease, is well defined [15] and can be diagnosed by clinical observations or by complementary examinations [16]. Use of screening tools (NRS 2002) and assessment (subjective global assessment (SGA), mini-nutrition assessment (MNA)) are recommended in the daily practice. A definition of acute critical illness-associated malnutrition still needs to be developed [8]. Meanwhile, following the screening process, the malnutrition assessment requires the association of a phenotype (weight loss %, BMI, decrease in

appetite, and/or low muscle mass) and an etiology e.g. critical illness [17] (GLIM (Global Leadership Initiative on Malnutrition) criteria).

Muscle mass can be evaluated by ultrasound [18], computerized tomography scan [19], or bioelectric impedance analysis (BIA) [20]. Sarcopenia, a decrease in muscle mass and/or function, is frequent in undernourished patients admitted to the ICU [21]. This loss of muscle may be considered as frailty and is associated with a prolonged hospital stay, and a decrease in quality of life and functional capacity [22]. Handgrip dynamometry can assess muscle function [23]. BIA can assess body composition in a stable patient not suffering from fluid compartment shifts and phase angle [24] is useful in the evaluation of the prognosis of critically ill patients. Patients with low muscle mass found at admission by computerized tomography have a higher length of stay and higher mortality [25].

5) To avoid overfeeding, early full EN and PN shall not be used in critically ill patients but shall be prescribed within three to seven days (Fig. 2).

(R8, Grade A, strong consensus, 100%)

Commentary

In a meta-analysis of studies comparing enteral and parenteral routes independent of timing, Elke et al. [26] found a reduction in ICU infections with EN as compared to PN (RR 0.64, 95% CI 0.48–0.87, $p = 0.004$, $I^2 = 47\%$). This difference did not occur when the amount of energy administered by PN and EN was similar (most recent studies), suggesting that caloric overfeeding may play a role in the infectious complications associated with PN and therefore in the decision process regarding the route, timing and the calorie target should also be taken into account. The energy/protein goal should be achieved progressively and not before the first 48–72 h to avoid over-nutrition. This progression should be ordered according to a local protocol preventing sharp and too rapid increases. Full targeted medical nutrition therapy is considered to achieve more than 70% of the resting energy expenditure (REE), but not to

exceed 100% of measured EE. Provision of excessive amounts of nutrients by any route should be avoided in the early phase of critical illness, which is associated with relevant endogenous energy production.

3.2. Medical nutrition therapy

3.2.1. Early medical nutrition (Fig. 3)

6) If oral intake is not possible, early EN (within 48 h) in critically ill adult patients should be performed/initiated rather than delaying EN.

(R4, Grade B, strong consensus, 100%)

Commentary

In comparing early EN with delayed EN (six studies in ICU patients and four studies including non-ICU patients), and similar to an earlier meta-analysis [13], a reduction of infectious complications with early EN (RR 0.76, CI 0.59, 0.97, $p < 0.03$) was observed [4]. However, this was true only when including studies that also enrolled patients outside of the ICU. There were no differences in other outcomes between early and delayed EN. Excluding earlier studies (before 2000) attenuates the signal that early EN may reduce infectious complications compared to delaying EN beyond 48 h. Importantly, the dosage of EN was not taken into consideration in this meta-analysis.

7) If oral intake is not possible, early EN (within 48 h) shall be performed/initiated in critically ill adult patients rather than early PN.

(R5, Grade A, strong consensus 100%)

Commentary

When comparing early EN with early PN (six studies in ICU patients and seven studies with non-ICU patients included) a reduction of infectious complications with EN (RR 0.50, 95%CI 0.37–0.67, $p = 0.005$), as well as shorter ICU (RR -0.73, 95%CI -1.30 to -0.16, $p = 0.01$) and hospital stay (RR -1.23, 95%CI -2.02 to -0.45, $p = 0.002$) was observed, while mortality was not different [4]. However, recent randomized controlled trials (RCTs) do not demonstrate a clear advantage of EN over PN, and the observed benefit of EN in earlier studies may be due to the higher energy and amino acid/protein content provided by PN compared to EN.

8) Early EN should be performed

- in patients receiving extracorporeal membrane oxygenation (ECMO)
- in patients with traumatic brain injury
- in patients with stroke (ischemic or hemorrhagic)
- in patients with spinal cord injury
- in patients with severe acute pancreatitis
- in patients after gastrointestinal surgery
- in patients after abdominal aortic surgery
- in patients with abdominal trauma when the continuity of the gastrointestinal tract is confirmed/restored
- in patients receiving neuromuscular blocking agents
- in patients managed in prone position
- In patients with open abdomen
- regardless of the presence of bowel sounds unless bowel ischemia or obstruction is suspected
- in patients with diarrhea

(R40, Grade B, strong consensus, 96%)

Commentary

The European Society of Intensive Care Medicine (ESICM) guidelines formulated 17 recommendations favoring initiation of early EN (within 48 h of ICU admission) and seven recommendations favoring delaying EN [13], as summarized in our recommendations No. 8, 11, 12. In meta-analyses performed for the ESICM guidelines, early EN reduced infectious complications in unselected critically ill patients, in patients with severe acute pancreatitis, and after gastrointestinal surgery, whereas no evidence of superiority for early PN or delayed EN over early EN was detected in any of the sub-questions. However, all issued recommendations were weak due to the low quality of evidence, with most of them finally based on expert opinion [13].

9) Early and progressive PN can be provided instead of no nutrition in case of contraindications for EN in severely malnourished patients.

(R7, Grade 0, strong consensus, 95%)

Commentary

EN is not feasible when a patient is determined to be at high nutrition risk (e.g., NRS 2002 ≥ 5) or severely malnourished, and, the initiation of low-dose PN should be carefully considered and balanced against the risks of overfeeding and refeeding, which may outweigh the expected benefits.

10) In patients who do not tolerate full dose EN during the first week in the ICU, the safety and benefits of initiating PN should be weighed on a case-by-case basis.

(R20, Grade GPP, strong consensus, 96%)

Commentary

There is no debate regarding the need for supplementing PN to EN in the case of prolonged nutritional deficit. However, the best timing to prescribe supplemental PN remains debated. Casaer et al. [14] observed that early (supplemental or exclusive) PN is associated with increased morbidity including prolonged ICU dependency and mechanical ventilation, and increased infection rate and need for renal replacement therapy. These findings may be related to the specific study protocol, the patients' characteristics and the large amounts of energy administered under guidance of predictive equations instead of indirect calorimetry. However, results of this study revealed the potential harm of nutritional intervention aiming at full, possibly overestimated calorie targets during the acute phase of critical illness. In addition, it is not known whether use of calorimetry would have resulted in different targets and different outcomes in the EPaNIC study. The optimal time point for supplemental PN aiming to achieve full caloric needs is not clear, but is suggested to be between days four and seven.

3.2.2. Delayed medical nutrition (Fig. 4)

11) EN should be delayed in several conditions displayed on Fig. 4

- if shock is uncontrolled and hemodynamic and tissue perfusion goals are not reached, whereas low dose EN can be started as soon as shock is controlled with fluids and




Fig. 4. Delayed medical nutrition (enteral/parenteral) in the ICU patient. Abbreviations: see Fig. 1.

vasopressors/inotropes, while remaining vigilant for signs of bowel ischemia;

- in case of uncontrolled life-threatening hypoxemia, hypercapnia or acidosis, whereas EN can be started in patients with stable hypoxemia, and compensated or permissive hypercapnia and acidosis;
- in patients suffering from active upper gastrointestinal bleeding, whereas EN can be started when the bleeding has stopped and no signs of re-bleeding are observed;
- in patients with overt bowel ischemia;
- in patients with high-output intestinal fistula if reliable feeding access distal to the fistula is not achievable;
- in patients with abdominal compartment syndrome; and
- if gastric aspirate volume is above 500 mL/6 h.

(R38, Grade B, strong consensus, 100%)

Commentary

This recommendation is based on a previous recommendation published by the ESCIM [13]. See commentary to No. 8.

12) Low dose EN should be administered

- in patients receiving therapeutic hypothermia and increasing the dose after rewarming;
- in patients with intra-abdominal hypertension without abdominal compartment syndrome, whereas temporary reduction or discontinuation of EN should be considered when intra-abdominal pressure values further increase under EN; and
- in patients with acute liver failure when acute, immediately life-threatening metabolic derangements are controlled with or without liver support strategies, independent on grade of encephalopathy.

(R39, Grade B, strong consensus, 96%)

Commentary

This recommendation is based on a previous recommendation published by the ESCIM [13]. See commentary to No. 8.

3.2.3. Enteral nutrition – route of administration (Fig. 4)

- 13) In patients deemed to be at high risk for aspiration, postpyloric, mainly jejunal feeding can be performed.**
(R12, Grade GPP, strong consensus, 95%)

Commentary

Patients with a very high risk of aspiration may benefit from early postpyloric EN. We recommend postpyloric feeding in patients with a high risk for aspiration. According to the American Society for Parenteral and Enteral Nutrition (ASPEN) recommendations [27], patients at increased risk for aspiration may be identified by a number of factors, including inability to protect the airways, support with mechanical ventilation, age >70 years, reduced level of consciousness, poor oral care, inadequate nurse:patient ratio, supine positioning, neurologic deficits, gastroesophageal reflux, transport out of the ICU, and use of bolus intermittent EN [28]. The Canadian Critical Care Practice Guideline guidelines [29] confirm this approach: "Strategies to Optimize Delivery and Minimize Risks of EN: Small Bowel Feeding vs. Gastric. Based on eleven level 2 studies, small bowel feeding compared to gastric feeding may be associated with a reduction in pneumonia in critically ill patients."

- 14) Continuous rather than bolus EN should be used (Fig. 4).**
(R9, Grade B, strong consensus, 95%)

Commentary

Five studies were identified and meta-analysis found a significant reduction in diarrhea with continuous versus bolus administration (RR 0.42, 95%CI 0.19–0.91, $p = 0.03$), whereas no difference was identified in other outcomes [4]. Despite the fact that bolus administration is significantly different from continuous feeding in healthy individuals, it significantly increases gastric volume and superior mesenteric artery blood volume in critically ill patients [30], although these differences are not always translated into clinical advantages. This limited amount of data suggests that bolus and continuous enteral feeding can achieve the same target without an increase in side effects in any of these routes. Finally, bolus feeding could provide a greater stimulus for protein synthesis [31].

15) Gastric access should be used as the standard approach to initiate EN.

(R10, Grade GPP, strong consensus, 100%) (Fig. 4)

Commentary

Our meta-analysis [4] shows that feeding intolerance was more prevalent in the case of gastric feeding in five studies (RR 0.16, 95% CI 0.06–0.45, $p = 0.0005$). We observed a trend for less pneumonia (eleven studies) (RR 0.75, 95%CI 0.55–1.03, $p = 0.07$) in patients treated with postpyloric feeding with no differences in mortality (twelve studies), diarrhea (seven studies) or ICU length of stay. As postpyloric tube placement requires expertise, is commonly associated with some time delay, and is considered less physiological compared to gastric EN, the routine use of the postpyloric route is currently not justified. Moreover, postpyloric feeding could possibly be harmful in cases of gastrointestinal motility problems distal to the stomach. Taken together, we suggest using gastric access as a standard and implementing postpyloric access in the case of intolerance to gastric feeding due to gastroparesis.

16) In patients with gastric feeding intolerance not solved with prokinetic agents, postpyloric feeding should be used (Fig. 4).

(R11, Grade B, strong consensus, 100%)

Commentary

Postpyloric EN has been associated with a decrease in ventilator acquired pneumonia in several earlier meta-analyses, but this benefit did not translate into decreases in length of ventilation, ICU or hospital stay, or mortality [4]. It should be considered that postpyloric tube placement requires expertise, is probably less physiologic compared to gastric EN, and could possibly be harmful in cases of GI motility problems distal to the stomach.

3.2.4. Parenteral nutrition (Fig. 4)

17) In case of contraindications to oral and EN, PN should be implemented within three to seven days.

(R6, Grade B, consensus 89%)

Commentary

When to start, which route to prefer and how to progress PN have been a matter of debate for years. Recent guidelines written by ESPEN [1,2], ASPEN/SCCM [32], the Canadian Critical Care Practice Guideline group [29] and the most recent clinical practice guidelines on early EN in critically ill patients by the ESICM working

group on gastrointestinal function [13] were considered when formulating the updated ESPEN recommendations [4]. The latter performed an extensive review of the literature, multiple meta-analyses, six web-seminars and utilized the GRADE methodology, evidence to decision framework and Delphi methodology. Since many of the authors of the current guidelines are also co-authors of the ESICM guidelines, a decision was made to endorse the respective recommendations related to early enteral feeding. Following the literature search we could agree with other guideline statements such as the ASPEN/SCCM guidelines [27] suggesting the "use of EN over PN in critically ill patients who require nutrition support therapy" (Evidence low to very low). The Canadian Critical Care Practice Guideline guidelines [29] recommend similarly stating "when considering nutrition support for critically ill patients, we recommend the use of EN over PN in patients with an intact gastrointestinal tract."

18) PN should not be started until all reasonable strategies to improve EN tolerance have been attempted.

(R 21, updated, Grade GPP, consensus, 100%)

Commentary

1. Revised indications/contraindications of PN considering dosage: Early full feeding independent of the route may be associated with impaired outcomes [33]. In fact, early full EN may even be more detrimental than early full PN [34].
2. Complications of EN and evidence on enteral feeding intolerance (EFI): EFI is defined as worsening gastrointestinal function in response to feeding attempts. Full EN may be associated with severe complications in shock patients [33,35]. Enhancing EN tolerance and reducing complications lacks a definitive strategy. Proposed interventions include:
 - a) Using adequate indications and contraindications for EN (according to recommendation 38)
 - b) Using postpyloric feeding in patients with gastric feeding intolerance/high risk for aspiration (according to recommendations 11 and 12), while being aware of the risk of bowel distension and mesenteric ischemia (very low evidence) [36]. Vomiting/gastric residual volume (GRV) monitoring may be needed to achieve safety.
 - c) Using prokinetic drugs as indicated to prevent complications, rather than for maximizing EN provision. Energy-dense feed delivery is associated with increased prokinetic use, but does not affect other outcomes [37]. A better survival was observed in patients in whom EFI was resolved within 72 h, more likely in patients receiving prokinetics [38].
 - d) Using adequate definitions for EN-associated complications to define EFI: abdominal distention, gastric overfilling, vomiting, diet regurgitation, bowel distension up to Ogilvie's syndrome and/or intestinal ischemia, diarrhea. Definition of EFI should not be limited solely to gastric function [39].
 - e) Considering intra-abdominal pressure to initiate and maintain, or not, EN [40].
 - f) Monitoring of gastrointestinal function with the use of scores such as the Gastrointestinal Dysfunction Score (GIDS) (not yet validated) [41].

3.2.5. Energy and protein targets (Fig. 5)

19) In critically ill mechanically ventilated patients, energy expenditure (EE) should be determined by using indirect calorimetry.

(R15, Grade B, strong consensus, 95%)

Commentary

The weakness of predictive equations and the use of indirect calorimetry have been subject to multiple evaluations and recommendations from ESPEN [2] and ASPEN [27], both preferring the use of indirect calorimetry to evaluate ICU patient actual energy expenditure. The predictive equations are associated with significant inaccuracy (up to 60%), leading to over or under evaluation of the needs and inducing over or underfeeding [42]. Numerous meta-analyses have demonstrated the poor value of predictive equations [43,44], variability that is increased because body weight remains difficult to accurately assess [45].

- 20) If indirect calorimetry is used, isocaloric nutrition rather than hypocaloric nutrition can be progressively implemented after the early phase of acute illness.**
(R16, Grade 0, strong consensus, 95%) (Fig. 5)

Commentary

Our meta-analysis focused only on studies using indirect calorimetry found a trend (RR 1.28, 95%CI 0.98–1.67, $p = 0.07$) to improved short-term mortality when using indirect calorimetry to identify the energy target, compared to hypocaloric regimens but there were no significant differences in long term mortality, infection or length of stay [4]. Four RCTs have based their energy targets on indirect calorimetry. The pilot TICACOS study [46] showed that such a strategy was associated with an improvement in 60 day survival in the per protocol study, but also to an increase in length of ventilation, infections and length of stay related to the calorie overload and positive energy balance due to non-nutritional energy intakes. Petros et al. [47] showed a reduction in infection rate in the study group. Heidegger et al. [48] measured EE at day 3 and adapted the energy intake accordingly, comparing supplemental PN from day four to an EN only group. The intervention group had a lower late nosocomial infection rate after day 9. The recent EAT-ICU study compared the goal-directed group, receiving

the EE measured with indirect calorimetry as a caloric target to reach within 24 h to patients receiving standard therapy. The study group also received protein according to urinary nitrogen loss. No advantages or harm were observed in terms of functional outcome, morbidity, or mortality in this RCT [49].

- 21) Statement: If calorimetry is not available, using VO_2 (oxygen consumption) from pulmonary arterial catheter or VCO_2 (carbon dioxide production) derived from the ventilator will give a better evaluation on EE than predictive equations.**
(S2, consensus, 82%)

Commentary

If indirect calorimetry is not available, calculation of REE from VCO_2 only obtained from ventilators ($REE = VCO_2 \times 8.19$) has been demonstrated to be more accurate than equations [50] but less accurate than indirect calorimetry [51]. VO_2 calculated from pulmonary artery catheter can also be used. In the absence of indirect calorimetry, VO_2 or VCO_2 measurements, the use of simple weight-based equations (such as 20–25 kcal/kg/d) may be preferred [1,2,27].

- 22) If predictive equations are used to estimate the energy need, hypocaloric nutrition (below 70% estimated needs) should be preferred over isocaloric nutrition for the first week of ICU stay.**
(R19, Grade B, strong consensus, 95%)

Commentary

Studies using predictive equations and observational studies were analyzed [4]. If predictive equations are used, we suggest using hypocaloric nutrition (up to 70% estimated needs), over isocaloric nutrition (70% or greater of estimated needs), in the early phase of acute illness (RR 0.92, 95%CI 0.86–0.99, $p = 0.02$). Various studies have compared energy intake based on predictive equations




Fig. 5. Energy and protein targets for ICU patients. EE, energy expenditure.

to reduced calorie intake achieving even trophic enteral feeding concluding that there was no difference between normocaloric versus hypocaloric diets in critically ill patients [4]. Berger & Pichard observe an increase in mortality in the group of patients receiving energy close to the prescribed recommended energy intake [52]. Conversely, Reignier et al. observed a better outcome in patients receiving low energy and proteins over 7 days (Nutrirea 3 = ref 33). Large observational series including hundreds to thousands of patients have observed that the optimal calorie load associated with the best survival is around 80% of predicted energy needs [53], whereas others suggested no relation between intake and outcome or better outcome with lower energy intakes [54]. However, in all these studies, calorie delivery was lower than recommended/prescribed or the studies were not targeted to this parameter.

23) Hypocaloric nutrition (not exceeding 70% of EE) should be administered in the early phase of acute illness (Fig. 5). (R17, Grade B, strong consensus, 100%)

Commentary

A larger database analysis suggested that energy intake is associated with significantly improved survival when it is close to measured EE [55] or between 70 and 100% of the repeatedly measured resting EE [56]. Undernutrition or over-nutrition is deleterious to outcome according to these large observational studies. If there is consensus stating that overfeeding should be avoided, it remains difficult to define which calorie targets should be proposed in the different phases of critical illness. Actual EE should not be the target during the first 72 h of acute critical illness. Early full feeding causes overfeeding as it adds to the endogenous energy production which amounts to 500–1400 kcal/d and can lead to deleterious effects such as increased length of stay, ventilation duration and infection rates [57]. Early full feeding also increases the risk of refeeding (see No. 60). On the other hand, a too low intake, below 50%, was associated in retrospective studies with a worse clinical outcome, may lead to severe calorie debt and empty the energy reserves, reduce lean body mass and may increase infectious complications [58,59]. However, prospective studies (Nutrirea-3 Permit EPANIC) comparing two levels of energy intake found a benefit associated with early provision of lower amounts.

24) After day three, energy delivery can be increased up to 80–100% of measured EE. (R18, Grade 0, strong consensus, 95%)

Commentary

Recently the analysis of a large data base including 1171 patients with indirect calorimetry data [56] confirmed that under- and overfeeding were both deleterious, and that the optimal amount appeared to be between 70 and 100% of measured EE. Prospective randomized studies comparing the delivery of 70–80% of the measured EE to another regimen may improve our knowledge.

25) During critical illness, 1.3 g/kg protein equivalents per day can be delivered progressively. (R22, updated, Grade 0, strong consensus, 92%)

Commentary

Numerous observational studies showed associations between high-dose protein delivery and improved outcomes

(morbidity or mortality). However, RCTs did not confirm the expected benefit of higher protein intakes alone, or combined with physical activity or energy [37,54,58]. Recently [60], a comparison between 1.6 ± 0.5 g/kg/d compared with 0.9 ± 0.3 g/kg/d administered protein did not find any significant difference in outcomes, but an increased mortality in patients with acute kidney injury randomized to high protein intakes. The recruited patients' energy intake was very variable, although similar in the groups (around 14 kcal/kg/d) receiving high and low-dose proteins. In accordance with the Lee et al. meta-analysis, there is no support to recommend higher doses than 1.3 g/kg/d, except for increased muscle mass in individual patients. Medium doses (not low and not high) are associated with the best outcomes [61]. The timing of administration has been mainly studied retrospectively or in post hoc analysis [62]. Recent stable isotope studies [63,64] demonstrate that amino acid conversion is improved with time, allowing an increase in whole body protein production only after the early period of the acute phase, even increasing further in the post-acute phase. This supports the progressive increase of protein intakes. There is a need for well-conducted RCTs to answer the question of the best timing and dose of protein administration in the ICU.

26) Statement 3: Physical activity may improve the beneficial effects of nutritional therapy. (S 3, consensus, 86%)

Commentary

Exercise has been suggested in several studies [65,66] to be effective in preventing anabolic resistance, reducing morbidity and improving the level of activity. Administration of increased protein intake together with increased physical activity during the late phase of critical illness should be further explored.

3.2.6. Other substrates (Fig. 6)

27) The amount of glucose (PN) or carbohydrates (EN) administered to ICU patients should not exceed 5 mg/kg/min. (R23, Grade GPP, strong consensus, 100%)

Commentary

The optimal nutritional composition of macronutrients is defined by minimal requirements and upper limits. Carbohydrates are the preferential substrate for production of energy, but in critical illness, insulin resistance and hyperglycemia are common secondary to stress. A minimal requirement has been proposed in previous guidelines [2] based on a society recommendation [67]. This evaluation is weak as has been stated: "carbohydrate could be theoretically eliminated from the diet, but it is probably safe® to give 150 g/d: This may be explained by organ dependence on glucose such as the brain (1–0–120 g/d), red blood cells, immune cells, renal medulla and all the transparent tissues of the eyes" [2] and the risks associated with hypoglycemia. The exact optimal carbohydrate amount to administer is difficult to determine. Endogenous glucose production is increased and does not decrease when nutrients and insulin are administered as compared with healthy conditions [68]. Excessive glucose based energy provision is associated with hyperglycemia, enhanced CO₂ production, enhanced lipogenesis, increased insulin requirements and no advantage in protein sparing in comparison with lipid based energy provision [69]. The use of diabetic-specific enteral

formula in ICU patients suffering from Type 2 Diabetes Mellitus decrease the requirement for insulin [70,71] and. The recommended glucose administration should not exceed 5 mg/kg/min [2].

- 28) The administration of intravenous lipid emulsions should be generally a part of PN.**
(R24, Grade GPP, strong consensus, 100%) (Fig. 6)

Commentary

Essential fatty acids (FA) were previously recommended at a dose of 8 g/d, but recent studies have shown that pediatric patients receiving pure fish oil lipid emulsions did not develop essential FA deficiency even after months. Lipid metabolism is modified in critical illness and low plasma triglyceride levels and high-density lipoprotein (HDL) cholesterol levels are associated with improved survival [72]. Administration of marked amounts of carbohydrates and lipids can lead to hyperglycemia and liver function test abnormalities while high fat administration can lead to lipid overload, and especially unsaturated fat to impaired lung function and immune suppression [73]. Close monitoring of triglycerides and liver function tests may guide the clinician [74].

- 29) Intravenous lipid (including non-nutritional lipid sources) should not exceed 1.5 g lipids/kg/d and should be adapted to individual tolerance.**
(R25, Grade GPP, strong consensus, 100%) (Fig. 6)

Commentary

For intravenous lipids, the upper recommendation is 1 g/kg body weight/d with a tolerance up to 1.5 g/kg/d. Administration in excess can lead to waste, storage or even toxicity.

Special attention should be paid if propofol is administered, since it is a source of FAs. This lipid solution contains 1.1 kcal/mL and can provide a large energy load exceeding nutritional support [75]. Electronic patient data management systems help to recognize this calorie overload. Citrate use in continuous veno-venous hemo-diafiltration is also associated with increased carbohydrate load and should be taken into account as a non-nutritional calorie intake [75].

Regarding the FA composition of the lipid emulsions, recent expert recommendations indicated that a blend of FAs should be considered, including medium chain triglycerides (MCTs), omega-9 mono-unsaturated FAs, and omega-3 polyunsaturated FAs. At this stage, the evidence for omega-3 FA-enriched emulsions in non-surgical ICU patients is not sufficient to recommend it as a standalone [76].

- 30) In patients with burns >20% body surface area, additional enteral doses of glutamine (GLN) (0.3–0.5 g/kg/d) should be administered for 10–15 days as soon as EN is commenced.**
(R26, Grade B, strong consensus, 95%)

Commentary

The amino acid GLN is a normal component of proteins, representing around 8% of all amino acids, and is present in standard




Fig. 6. Nutritional substrates other than protein needed for treatment of ICU patients.

commercial enteral feeds. GLN for parenteral use has been available since 1994, after its synthesis by Fürst and Stehle [77]. For stability reasons, it was not present in standard PN.

In major burns, studies include a limited number of patients: nevertheless, the existing randomized trials have repeatedly demonstrated that GLN (and its precursor ornithine α -ketoglutarate) has beneficial effects in major burn injuries, reducing infectious complications (mainly gram negative infections) and also mortality [4]. This has been confirmed in the latest meta-analysis [4], and is included in the specific ESPEN burn guidelines [78], but is challenged by the largest RCT revealing no benefit from GLN supplementation in burns [79]. A higher GLN requirement in burn patients is explained by exudative losses: analysis of burn exudates shows that GLN is lost in larger amounts than any other amino acid [80].

- 31) In critically ill trauma, additional EN doses of GLN (0.2–0.3 g/kg/d) can be administered for the first five days with EN. In case of complicated wound healing, it can be administered for a longer period of ten to 15 days (Fig. 6). (R27, Grade 0, strong consensus, 91%)**

Commentary

The efficiency of enteral GLN on infection reduction was also suggested in major trauma. A RCT in 20 trauma patients with delayed wound healing, showed that oral antioxidant and GLN containing supplements reduced time to wound closure (22 days versus 35; $p = 0.01$). In the control patients a decline of plasma GLN was observed, while it was modestly increased in those having received 20 g GLN per day for 14 days. Finally, enteral GLN was shown to improve body composition and in particular lean body mass in a group of 44 head and neck cancer patients randomized to receive a GLN supplement (30 g daily) for four weeks [81]. The authors observed a significant improvement of fat-free mass, serum albumin, and quality of life scores post-operatively [81].

- 32) In ICU patients except burn and trauma patients, additional enteral GLN should not be administered (Fig. 6). (R28, Grade B, strong consensus, 92%)**

Commentary

Rodas et al. [82] showed a U-shaped association between plasma GLN levels and outcome.

In other critically ill patients, the MetaPlus trial [83] showed no advantage in terms of infection of a feeding solution containing additional enteral GLN. Of note, none of the groups received the planned high dose protein resulting in a mean delivery of 0.9 g/kg/d.

- 33) In unstable and complex ICU patients, particularly in those suffering from liver and renal failure, parenteral GLN-dipeptide shall not be administered (Fig. 6). (R29, Grade A, strong consensus, 92%)**

Commentary

Since the 1990s, many studies have shown benefits in terms of infectious complication reduction, lower mortality and reduction of hospital costs recently confirmed in an analysis including RCTs performed after 2000, using GLN as part of nutrition support [84].

When analyzed together [84] most single center studies observed improved survival while some multicenter studies did not confirm this finding. The positive trials used GLN as part of global

nutrition in stabilized patients. The REDOXs study [85], designed as a 2 x 2 factorial trial, generated concerns for several reasons, including the fact that the randomization was associated with higher severity with more organ failures in the GLN groups upon enrollment, largely explaining the higher mortality. Finally, Stehle et al. [86] in a meta-analysis including only stable patients showed an advantage to administering GLN.

- 34) High doses of omega-3-enriched EN formula should not be given by bolus administration. (R30, Grade B, strong consensus, 91%)**

Commentary

Aggregating all the studies in ICU patients without taking into account the amount of omega-3 FAs or whether they are given as bolus or continuous administration, does not yield any advantage for any formula [87]. Glenn and Wischmeyer [88] analyzed separately the studies administering omega-3 FAs as a bolus or in a continuous manner and found that continuous administration improved length of stay and length of ventilation; in contrast, bolus administration had no advantage. The pre-emptive administration of the same formula administered in three studies in severe, ventilated, multiple trauma patients did not find any advantage [89]. In at least one study, the membrane content of eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA) was very low at baseline and was hardly corrected with omega-3 and borage oil administration, suggesting that we do not know the exact amount of omega-3 FAs to administer to this category of patients.

- 35) EN enriched with omega-3 FA within nutritional doses can be administered. (R31, Grade 0, strong consensus, 95%)**

Commentary

Eight studies addressing this question were identified; in four of them antioxidants were also given. A meta-analysis did not reveal any benefit of omega-3 FA, but there was a trend towards increase in PO_2/FiO_2 with intervention (RR 22.59, 95%CI -0.88 - 46.05, $p = 0.06$). However, because it may change quickly and is dependent on ventilator settings, fluid status, body position etc. PO_2/FiO_2 is probably not the best outcome variable [4].

Enteral formulae enriched in borage oil and/or omega-3 FAs have been administered in patients suffering from adult respiratory distress syndrome, acute lung injury and sepsis with positive effects regarding length of stay, length of ventilation and even mortality [4]. Our meta-analysis found no significant advantage in oxygenation for enteral formulae enriched in omega-3 FAs (mainly EPA), gamma-linolenic acid and antioxidants over control groups receiving lipid-rich formulas [4].

- 36) High doses omega-3 enriched enteral formulas should not be given on a routine basis. (R32, Grade B, consensus, 90%)**

Commentary

In the post-hoc analysis of the MetaPlus study [90], administering GLN, EPA/DHA and antioxidants to critically ill patients, only the change from baseline to day 4 of EPA + DHA/long chain triglyceride (LCT) ratio was statistically significantly associated with six month mortality (hazard ratio 1.18, 95%CI 1.02–1.35, $p = 0.021$) suggesting a harmful effect of these nutrients in medical ICU patients. It has to be noted that this harmful effect was not observed in

the previous studies on patients in acute lung injury or adult respiratory distress syndrome.

- 37) Parenteral lipid emulsions enriched with EPA + DHA (Fish oil dose 0.1–0.2 g/kg/d) can be provided in patients receiving PN (Fig. 6).**
(R33, updated, Grade 0, strong consensus, 100%)

Commentary

Recently, several studies have shown the biological effects, clinical benefits, and safety of lipid emulsions with reduced content of 18-carbon omega-6 FA. The use of intravenous fat emulsions based solely on soybean oil (rich in 18-carbon omega-6 FA) should be avoided due to their likely pro-inflammatory effects [2]. Alternative lipid emulsions including sources incorporating olive oil, fish oil, and coconut oil (MCTs) in various combinations [76] are available. Mechanistic understandings of the action of omega-3 FA from fish oil have advanced [91], and positive effects of fish oil on organ function (kidney, liver, muscle) have been reported [92]. New trials and meta-analyses of fish oil containing PN in critically ill and surgical patients have been published [93]. Some meta-analyses combine trials of enteral and parenteral fish oil in critically ill patients [94,95]. Each of these meta-analyses favors fish oil for clinical benefit. Pradelli et al. [96] conducted a meta-analysis of 24 RCTs comparing fish oil-enriched PN with standard (non-fish oil-enriched) PN in adult patients in the ICU. In critically ill ICU patients, fish oil was associated with reductions in infections (RR 0.65, 95% CI 0.46–0.94), ICU LOS (–2.14 days, 95% CI –3.89 to –0.40), and hospital length of stay (–3.98 days, 95% CI –6.90 to –1.06), but with a non-significant reduction in 30-day mortality in all ICU patients (RR 0.90, 95% CI 0.69–1.16). It was estimated in cost-effectiveness analyses that parenteral fish oil would lower overall hospital costs (compared with standard PN) in all six countries considered.

- 38) To enable substrate metabolism, micronutrients (i.e. trace elements and vitamins) should be provided daily with PN (Fig. 6).**
(R34, Grade B, strong consensus, 100%)

Commentary

Providing micronutrients to include the full range of trace elements and vitamins is an integral part of nutritional support as stated in the 2009 guidelines [2]. Parenteral and enteral feeding preparations differ in that commercially available PN solutions contain no micronutrients for stability reasons: this requires their separate prescription [2]. There are no PN studies with or without micronutrients.

The blood levels of several micronutrients are below normal ranges during the inflammatory response, and hence the impact of their provision is difficult to interpret. Recent evidence tends to show that persistently low zinc concentrations might become an important biomarker of adverse outcomes in sepsis [97].

We recommend the repletion of micronutrients in conditions of chronic and acute deficiency. Continuous renal replacement therapy for more than two weeks is a new cause of acute micronutrient deficiencies and particularly of severe copper deficiency that may explain life-threatening complications in patients requiring this therapy [98].

- 39) Antioxidants as high-dose monotherapy or combination therapy should not be administered without proven**

deficiency (Fig. 6).
(R35, updated, Grade A, strong consensus 100%)

Commentary

High-quality RCTs evaluating high doses of selenium [99], vitamin D [100,101], vitamin C [102–108], and combinations of vitamin C and thiamine [109], revealed no hard clinical benefit of such interventions.

- 40) Vitamin D (25(OH)D) status can be determined in all patients considered at risk of vitamin D depletion or deficiency (Fig. 6).**
(R36, updated, Grade GPP, strong consensus 100%)

Commentary

Vitamin D3 can be synthesized in sufficient amounts by the human body so long as there is exposure to sunlight and good liver and renal function. Vitamin D3 has a nuclear receptor and a large number of genes are under direct or indirect control of this vitamin. Hypovitaminosis D is common in the general population, with a seasonal occurrence, while low plasma concentrations of vitamin D have been repeatedly shown in critically ill patients. In the latter patients, deficiency has been associated with poor outcome including excess mortality, longer length of stay, higher sepsis incidence, and longer mechanical ventilation [110]. There is an uncertainty regarding dosing and timing of vitamin D administration. Therefore, this recommendation replaces the 2 recommendations (R36 and R37) related to dosage in the previous guidelines.

3.3. Medical nutrition in special cases (Fig. 7)

3.3.1. The non-intubated patient

- 41) In non-intubated patients not reaching the energy target with an oral diet, oral nutritional supplements should be considered first and then EN.**
(R4 1, Grade GPP, strong consensus, 96%)

Commentary

Reeves et al. [111] described the energy and protein intakes of patients with adult respiratory distress syndrome receiving non-invasive ventilation. From this small observational study, it is concluded that oral intake was inadequate, mainly with increasing time on non-invasive ventilation, and earlier during their hospital admission. In total, 78% of the patients met less than 80% of the requirements. The authors recommended referring the patients recognized to have swallowing issues for swallowing evaluation, to prevent oral nutrition complications.

- 42) In non-intubated patients with dysphagia, texture-adapted food can be considered. If swallowing is proven unsafe, EN should be administered.**
(R4 2, Grade GPP, strong consensus, 94%)

Commentary

Oral intake is impaired **after extubation** and a high incidence of swallowing dysfunction has been described (between 10 and 67.5%, with a mean around 50%, despite different timing and methods used for assessing the dysphagia) [112]. This post-

extubation swallowing disorder could be prolonged for up to 21 days mainly in the elderly and after prolonged intubation. Thus, at 21 days post-extubation, 24% of older patients were feeding tube dependent [113]. Recently, 29% of 446 ICU patients had prolonged post-extubation swallowing disorder at discharge and some post-extubation swallowing disorder has been shown four months after discharge [114]. The same authors who described the tools to diagnose post-extubation swallowing disorder, also suggest the use of thickening food to increase oral intake.

43) In non-intubated patients with dysphagia and a very high aspiration risk, postpyloric EN or, if not possible, temporary PN during swallowing training with removed nasogastric tube can be performed.

(R4 3, Grade GPP, strong consensus, 92%)

Commentary

Oral intake is frequently prescribed in the intensive care setting varying from 25 to 45% of the patients in the first four days, but does not reach the energy or protein requirements according to the Nutrition Day ICU survey [6]. This population includes patients admitted for monitoring, patients receiving non-invasive ventilation and post intubation/tracheostomy patients.

After tracheostomy, a cohort study showed that the majority of the patients returned to oral intake, but the time to commencement of oral intake was correlated with increased time to decannulation and increased time to decannulation

correlated with increased hospital length of stay [115]. Supplemental PN has not been extensively studied in this population.

3.3.2. The surgical patient (Fig. 7)

44) In patients after abdominal or esophageal surgery, early EN can be preferred over delayed EN.

(R4 5, Grade 0, strong consensus, 96%)

Commentary

We performed a meta-analysis of EN vs no nutrition within the first 48 h, which did not reveal clear benefit of EN in this subgroup of patients, but a trend towards fewer infectious complications was observed (RR 0.47, 95%CI 0.20–1.07, $p = 0.07$) [4]. Importantly, the presence of an intestinal anastomosis or re-anastomosis without leakage should not delay EN.

45) In critically ill patients with surgical complications after abdominal or esophageal surgery and unable to eat orally, EN (rather than PN) should be preferred unless discontinuity or obstruction of gastrointestinal tract, or abdominal compartment syndrome is present (Fig. 7).

(R4 6, Grade GPP, strong consensus, 96%)

Commentary

Esophageal surgery commonly results in the loss of the lower esophageal sphincter function and is therefore associated with a




Fig. 7. Medical nutrition in special cases. Abbreviations: see Fig. 1.




Fig. 8. Monitoring of medical nutrition in the ICU. Abbreviations: see Fig. 1.

significantly increased risk of aspiration. Therefore, many centers use “nil per mouth” strategy with EN via a surgical jejunostomy. We identified two RCTs addressing early EN via surgical jejunostomy in patients after esophageal surgery (in one case, the study group included other upper gastrointestinal surgery patients, not limited to esophageal surgery), suggesting potentially beneficial effects on the inflammatory state when compared with early PN and lower infection rates when compared with delayed EN. One larger retrospective study comparing early EN via surgical jejunostomy with early PN resulted in less life-threatening complications and a shorter postoperative hospital stay [116].

- 46) In the case of an unrepaired anastomotic leak, internal or external fistula, a feeding access distal to the defect should be aimed for to administer EN.**
(R4 7, Grade GPP, strong consensus, 96%)

Without evidence, but based on common reasoning and pathophysiological considerations, surgical complications leading to gastrointestinal contents leaking into the abdominal cavity should always lead to withholding/stopping EN. At the time of developing such complications, patients usually have developed considerable energy deficits. Therefore, PN should be considered early after re-surgery if such a problem clearly cannot be solved within the next days, but started at a slow infusion rate. Enteral feeding access distal to the leak should be aimed for in these cases. Small bowel ischemia associated with early (in some cases aggressive) EN via surgical jejunostomy has been reported in several case reports [117,118]. In these cases, close monitoring of abdominal symptoms is required, and only continuous administration and slow build-up of EN via jejunostomy is advocated.

- 47) In the case of an unrepaired anastomotic leak, internal or external fistula, or if distal feeding access is not achieved, EN should be withheld and PN may be commenced.**
(R4 8, Grade GPP, strong consensus, 100%)

Commentary

Two studies addressing early EN vs early PN in elective upper gastrointestinal surgery were identified [119–121]. In a sub-group analysis of the EPaNIC study, early and late PN were compared in complicated pulmonary/esophageal and abdomino-pelvic surgery patients. Reduced infection rates in late vs early PN were observed (29.9% vs. 40.2%, $p = 0.01$) with no difference in any mortality outcomes, whereas all these patients received virtually no EN during the seven study days [14]. The latter finding should most likely be interpreted as a harmful effect of early full feeding, also demonstrated in several other recent studies.

- 48) In case of high output stoma or fistula, the appropriateness of chyme reinfusion or enteroclysis should be evaluated and performed if adequate (Fig. 7).**
(R4 9, Grade GPP, strong consensus, 100%)

Commentary

In many cases of complicated abdominal surgery, patient tolerance to EN is impaired. Furthermore, depending on surgery, mal-digestion and/or malabsorption may occur. Therefore (supplemental), PN should be considered timely to avoid prolonged nutritional deficits. In specific situations with high-output stoma or fistula, chyme reinfusion or entero/fistuloclysis should be considered [122].

3.3.3. The trauma patient (Fig. 7)

- 49) Trauma patients should preferentially receive early EN instead of early PN.**
(R 50, Grade B, strong consensus, 96%)

Commentary

Our meta-analysis including three studies showed a decrease in length of stay (RR -0.47, 95%CI -7.57 to -1.71, $p = 0.002$), and a trend for decrease in mortality (RR 0.69, 95%CI 0.39–1.23, $p = 0.21$), but no difference in incidence of pneumonia when early EN was administered rather than early PN [4].

Most trauma patients are not malnourished on admission (6% SGA C), but may become malnourished during ICU stay (increase in SGA B) [123]. Kompan et al. [124] compared early EN to early PN followed by EN in multiple trauma patients and found a significant decrease in pneumonia and length of stay, but not in hospital stay and mortality. Fan et al. [125] compared three groups: early EN, early PN and EN followed by supplemental PN. Complications and mortality were significantly decreased and nutritional status and clinical outcomes were improved in the early EN + supplemental PN group. An earlier meta-analysis [126] showed that early EN was associated with reduced mortality. Higher protein intake reaching 1.5–2 g/kg/d may be considered in this population, since there are large protein losses (20–30 g/L of abdominal fluid) [127].

3.3.4. The septic patient (Fig. 7)

- 50) Early and progressive EN should be used in septic patients after hemodynamic stabilization.**

If contraindicated, EN should be replaced or supplemented by progressive PN.
(R4 4, Grade GPP, strong consensus 94%)

Commentary

Elke et al. [128] showed that not receiving nutrition support is deleterious in 2270 patients with sepsis, pneumonia and with an ICU stay of more than three days. Increased amounts of calories and protein per day were associated with a decrease in 60-day mortality and an increase in ventilation-free days. Early versus late EN initiation were not different [129]. Hypocaloric or trophic EN versus full EN showed no survival differences either [130]. To prevent a full provision of energy, a pragmatic approach remains to consider EN as the first choice for nutrition support during the first three to four days after ICU admission. If this is not feasible or is insufficient after three days, PN should be prescribed up to approximatively half of the predicted or measured energy needs and EN prescribed as soon as the clinical condition permits. Weijs et al. reported that septic patients did not improve outcome when receiving increased (1.2 g/kg/d) protein intake compared to non-septic patients [55].

Septic shock: Impaired splanchnic perfusion can potentially be further aggravated by EN administration and lead to bowel ischemia [131]. EN should be started after successful resuscitation [132]. In patients with sepsis, a fraction (20–50%) of full nutrition support should be initiated as early as possible to “open” the enteral route; then the amounts of feeds should be progressively increased according to the gastrointestinal tolerance in order to achieve optimal nutrition support once patients improved. If not feasible for prolonged periods, PN should be prescribed.

3.3.5. The obese patient (Fig. 7)

- 51) An iso-caloric high protein diet can be administered to obese patients, preferentially guided by indirect calorimetry measurements and urinary nitrogen losses.**
(R5 1, Grade 0, consensus, 89%)

Commentary

Overweight and obese patients have become more prevalent in ICUs. Reported recommendations [27] are summarized by Dickerson et al. [127]. There is a large variability in the prevalence between countries based on data from the Nutrition Day project [7]. If hypocaloric medical nutrition therapy appears to be the rule on many ICUs [7], we recommend the measurement of energy expenditure with indirect calorimetry and urinary nitrogen loss to guide energy requirements and protein needs, since predictive equations are inaccurate. Obese patients defined on the basis of BMI are a heterogeneous group. High BMI may be associated with an extremely trained muscle mass as in body builder at one end of the spectrum and sarcopenic obesity with an even lower muscle mass than would be expected from height at the other end. The muscle mass of obese patients will be highly dependent on their level of activity. Age is a further factor to be considered. Muscle mass typically is maximal between 25 and 35 years of age and decreases thereafter. Thus, in an older person with the same body weight, a lower muscle mass is likely to be present (see statement #4 for further details).

- 52) In obese patients, energy intake should be guided by indirect calorimetry.**

Protein delivery should be guided by urinary nitrogen losses or lean body mass determination (using computerized tomography or other tools).

If indirect calorimetry is not available, energy intake can be based on “adjusted body weight”.

If urinary nitrogen losses or lean body mass determination are not available, protein intake can be 1.3 g/kg “adjusted body weight”/d.

(R5 2, Grade GPP, consensus, 89%)

Commentary

If indirect calorimetry is not available and nitrogen excretion not measured, we suggest the use of ideal body weight as reference weight in overweight and obese patients. We propose to decrease energy provision where BMI indicates overweight or obesity. The reference (adjusted) body weight should then change from actual body weight to ideal body weight at a BMI >25 kg/m². Probably using as ideal body weight: 0.9 x height in cm –100 (male) (or –106 (female)) is sufficiently precise giving the overall uncertainties. Such an approach would completely ignore the metabolic demand of adipose tissue and muscle. Adipose tissue utilizes 4.5 kcal/kg/d and muscle 13 kcal/kg/d [133]. The proportion of muscle within the excess weight of an obese individual might be roughly 10%. A pragmatic approach is to add 20–25% of the excess weight (actual body weight-ideal body weight) to ideal body weight for all calculations of energy requirements.

Several authors advocate a controlled undernutrition of obese subjects while providing a relatively larger dose of protein between 2 and 2.5 g/kg/d (ideal body weight as reference) [134].

Additional metabolic derangements such as decreased glucose tolerance, altered lipid metabolism, lack of micronutrients and decreased gut motility will need specific attention [135]. Recommendations on early EN, gastrointestinal tolerance and progressive

increase in nutrition over several days apply similarly to overweight and obese patients as to all other ICU patients.

3.4. Monitoring of medical nutrition (Fig. 8)

3.4.1. Monitor for tolerance, overfeeding and underfeeding (see also rec. 5)

- 53) Blood glucose should be measured initially (after ICU admission or after artificial nutrition initiation) and at least every 4 h, for the first two days in general.**
(R5 3, Grade GPP, strong consensus, 93%)

Commentary

A number of observational studies confirmed a strong association between severe hyperglycemia (>180 mg/dL, 10 mmol/L) [136], marked glycemic variability (coefficient of variation >20%) [137], mild hypoglycemia (<70 mg/dL, 3.9 mmol/L) [138] and increased mortality. However, the prospective trials remain inconclusive, owing to differences in practices and to the difficulties in achieving safe and effective glycemic control. The glycemic target associated with the best adjusted outcome ranges from 80 to 150 to 140–180 mg/dL (7.8–10 mmol/L), which is different from the blood glucose levels actually achieved [139].

- 54) Insulin shall be administered, when glucose levels exceed 10 mmol/L.**
(R5 4, Grade A, strong consensus, 93%)

Commentary

Blood glucose control is essential and should be targeted. Current recommendations suggest starting insulin therapy when blood glucose exceeds 150 [43] or 180 mg/dL (10 mmol/L) [140].

In unstable patients even more frequent measurements may be required, whereas frequency can usually be decreased when a stable phase is reached, usually after 48 h.

The assessment and control of glycemia encompasses multiple steps [141]: Blood draw, use of blood gas analyzer or central laboratory analyzers (hexokinase-based) is essential, insulin: intravenous and continuous in case of ongoing nutrition support (enteral or parenteral) using an electric syringe and insulin algorithm.

- 55) In critically ill patients with gastric feeding intolerance, intravenous erythromycin should be used as a first line prokinetic therapy.**
(R13, Grade B, strong consensus, 100%)

Commentary

According to our meta-analysis [4] prokinetic use is associated with a trend towards better enteral feeding tolerance (RR 0.65, 95% CI 0.37–1.14, $p = 0.14$). This is significant for intravenous erythromycin (usually at dosages of 100–250 mg three times a day) (RR 0.58, 95%CI 0.34–0.98, $p = 0.04$) for two to four days but not for other prokinetics like metoclopramide (at usual doses of 10 mg two to three times a day). Effectiveness of erythromycin or other prokinetics is decreased to one third after 72 h [142] and they should be discontinued after three days. Our meta-analysis based on six studies finds a significant advantage to erythromycin and its use should be encouraged for 24–48 h, since it promotes gastric motility, and if a large (>500 mL) GRV still persists, the use of post-

pyloric feeding should be considered over withholding EN, unless a new abdominal complication (obstruction, perforation, severe distension ...) is suspected [4].

- 56) Alternatively, intravenous metoclopramide or a combination of metoclopramide and erythromycin can be used as a prokinetic therapy.**
(R14, Grade 0, strong consensus, 100%)

Commentary

The measurement of GRV for the **assessment** of gastrointestinal dysfunction is common and may help to identify intolerance to EN during initiation and progression of EN. However, **monitoring** of established EN with continued measurements of GRV may not be necessary [143]. We suggest that enteral feeding should be delayed when GRV is > 500 mL/6 h. In this situation, and if examination of the abdomen does not suggest an acute abdominal complication, application of prokinetics should be considered. ASPEN/SCCM [27] and the Surviving Sepsis initiative [144] recommend the use of prokinetics metoclopramide (10 mg three times a day) and erythromycin (3–7 mg/kg/d) in the case of feeding intolerance (weak recommendation, low quality of evidence for the surviving sepsis initiative, and for ASPEN/SCCM) [27].

3.4.2. Monitor for refeeding

- 57) Electrolytes (potassium, magnesium, phosphate) should be measured at least once daily for the first week.**
(R5 5, Grade GPP, strong consensus, 92%).

Commentary

Refeeding syndrome can be defined as the potentially fatal shifts in fluids and electrolytes that may occur in malnourished patients receiving artificial refeeding. Each case of refeeding syndrome – a potentially lethal state [145] – has to be detected early to prevent complications [146]. Therefore, assessment of nutritional status at admission is needed with a schedule for the measurement of electrolytes, including phosphate. Laboratory parameters (electrolytes monitoring, glycemia, liver function tests) are important to prevent or detect severe complications like refeeding syndrome or liver dysfunction related to nutrition, as well as to assist in the achievement of normoglycemia and normal electrolyte values.

- 58) In patients with refeeding hypophosphatemia (<0.65 mmol/L or a drop of >0.16 mmol/L), electrolytes should be measured 2–3 times a day and supplemented if needed.**
(R56, Grade GPP, strong consensus, 100%)

Commentary

Repeated measurements of P, K and Mg during initiation of feeding in critically ill patients are important to detect development of refeeding syndrome, especially because among critically ill patients' electrolyte disturbances upon refeeding are not limited to patients with overt malnutrition.

- 59) In patients with refeeding hypophosphatemia energy supply should be restricted for 48 h and then gradually increased.**
(R57, Grade B, strong consensus, 100%)

Commentary

The occurrence of refeeding hypophosphatemia may be perceived as a warning signal. In a RCT, Doig et al. showed that protocolized caloric restriction for 48 h in patients developing hypophosphatemia upon refeeding improved survival despite similar phosphate supplementation in both groups [147]. Slow progressions to energy target during the first 72 h, also called caloric restriction, should be considered to facilitate control of electrolyte disturbances if refeeding syndrome is anticipated or detected [148]. Importantly, whereas potassium is commonly measured in critically ill patients, measurements of phosphate are less common. Undetected rapid development of severe hypophosphatemia may lead to death after initiation of feeding as patients admitted to ICU are often malnourished either before or during admission to the hospital. A recent early calorie restriction study showed that electrolyte alterations were less likely to occur with a cautious introduction of feeding and this was confirmed by a retrospective study [149].

Acknowledgement

We thank Simon JW Oczkowski and Waleed Alhazzani from the Dept. of Medicine, Division of Critical Care and Department of Clinical Epidemiology and Biostatistics, McMaster University, Hamilton, Canada, for the valuable contributions to the original ESPEN ICU guideline.

References

- [1] Kreymann KG, Berger MM, Deutz NEP, Hiesmayr M, Jolliet P, Kazandjiev G, et al. ESPEN guidelines on enteral nutrition: intensive care. *Clin Nutr* 2006;25:210–23.
- [2] Singer P, Berger MM, Van den Berghe G, Biolo G, Calder P, Forbes A, et al. ESPEN guidelines on parenteral nutrition: intensive care. *Clin Nutr* 2009;28:387–400.
- [3] Bischoff SC, Singer P, Koller M, Barazzoni R, Cederholm T, van Gossum A. Standard operating procedures for ESPEN guidelines and consensus papers. *Clin Nutr* 2015;34:1043–51.
- [4] Singer P, Blaser AR, Berger MM, Alhazzani W, Calder PC, Casaer MP, et al. ESPEN guideline on clinical nutrition in the intensive care unit. *Clin Nutr* 2019;38:48–79.
- [5] Singer P, Weinberger H, Tadmor B. Which nutritional regimen for the comorbid complex intensive care unit patient? *World Rev Nutr Diet: S. KARGER AG* 2012:169–78.
- [6] Wischmeyer PE. Tailoring nutrition therapy to illness and recovery. *Crit Care* 2017;21:316.
- [7] Bendavid I, Singer P, Theilla M, Themessl-Huber M, Sulz I, Mouhieddine M, et al. NutritionDay ICU: a 7 year worldwide prevalence study of nutrition practice in intensive care. *Clin Nutr* 2017;36:1122–9.
- [8] Kondrup J. Nutritional risk screening (NRS 2002): a new method based on an analysis of controlled clinical trials. *Clin Nutr* 2003;22:321–36.
- [9] Ellia M. The 'MUST' report. Nutritional screening for adults: a multidisciplinary responsibility. *Redditch: BAPEN*; 2003.
- [10] Heyland DK, Dhaliwal R, Jiang X, Day AG. Identifying critically ill patients who benefit the most from nutrition therapy: the development and initial validation of a novel risk assessment tool. *Crit Care* 2011;15:R268 [R].
- [11] Arabi YM, Preiser J-C. A critical view on primary and secondary outcome measures in nutrition trials. *Intensive Care Med* 2017;43:1875–7.
- [12] Lew CCH, Yandell R, Fraser RJL, Chua AP, Chong MFF, Miller M. Association between malnutrition and clinical outcomes in the intensive care unit: a systematic review. *J Parenter Enteral Nutr* 2016;41:744–58.
- [13] Reintam Blaser A, Starkopf J, Alhazzani W, Berger MM, Casaer MP, Deane AM, et al. Early enteral nutrition in critically ill patients: ESICM clinical practice guidelines. *Intensive Care Med* 2017;43:380–98.
- [14] Casaer MP, Mesotten D, Hermans G, Wouters PJ, Schetz M, Meyfroidt G, et al. Early versus late parenteral nutrition in critically ill adults. *N Engl J Med* 2011;365:506–17.
- [15] Fried LP, Tangen CM, Walston J, Newman AB, Hirsch C, Gottdiener J, et al. Frailty in older adults: evidence for a phenotype. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences* 2001;56:M146–57.
- [16] Cederholm T, Bosaeus I, Barazzoni R, Bauer J, Van Gossum A, Klek S, et al. Diagnostic criteria for malnutrition – an ESPEN consensus statement. *Clin Nutr* 2015;34:335–40.
- [17] Cederholm T, Jensen GL, Correia M, Gonzalez MC, Fukushima R, Higashiguchi T, et al. GLIM criteria for the diagnosis of malnutrition – a consensus report from the global clinical nutrition community. *Journal of cachexia, sarcopenia and muscle* 2019;10:207–17.
- [18] Wischmeyer PE, San-Millan I. Winning the war against ICU-acquired weakness: new innovations in nutrition and exercise physiology. *Crit Care* 2015;19(Suppl 3):S6 [S].
- [19] Looijgaard WGP, Dekker IM, Stapel SN, Girbes ARJ, Twisk JWR, Oudemans-van Straaten HM, et al. Skeletal muscle quality as assessed by CT-derived skeletal muscle density is associated with 6-month mortality in mechanically ventilated critically ill patients. *Crit Care* 2016;20:386.
- [20] Thibault R, Makhoulouf A-M, Mulliez A, Cristina Gonzalez M, Kekstas G, Kozjek NR, et al. Fat-free mass at admission predicts 28-day mortality in intensive care unit patients: the international prospective observational study Phase Angle Project. *Intensive Care Med* 2016;42:1445–53.
- [21] Studenski SA, Peters KW, Alley DE, Cawthon PM, McLean RR, Harris TB, et al. The FNIH sarcopenia project: rationale, study description, conference recommendations, and final estimates. *J Gerontol A Biol Sci Med Sci* 2014;69:547–58.
- [22] Detsky AS, Baker JP, Mendelson RA, Wolman SL, Wesson DE, Jeejeebhoy KN. Evaluating the accuracy of nutritional assessment techniques applied to hospitalized patients: methodology and comparisons. *J Parenter Enteral Nutr* 1984;8:153–9.
- [23] Fan E, Ciesla ND, Truong AD, Bhoopathi V, Zeger SL, Needham DM. Inter-rater reliability of manual muscle strength testing in ICU survivors and simulated patients. *Intensive Care Med* 2010;36:1038–43.
- [24] Reis de Lima e Silva R, Porto Sabino Pinho C, Galvão Rodrigues I, Gildo de Moura Monteiro Júnior J. Phase angle as an indicator of nutritional status and prognosis in critically ill patients. *Nutr Hosp* 2014;31:1278–85.
- [25] Braunschweig CA, Sheehan PM, Peterson SJ, Gomez Perez S, Freels S, Troy KL, et al. Exploitation of diagnostic computed tomography scans to assess the impact of nutrition support on body composition changes in respiratory failure patients. *JPEN - J Parenter Enter Nutr* 2014;38:880–5.
- [26] Elke G, van Zanten ARH, Lemieux M, McCall M, Jeejeebhoy KN, Kott M, et al. Enteral versus parenteral nutrition in critically ill patients: an updated systematic review and meta-analysis of randomized controlled trials. *Crit Care* 2016;20:117.
- [27] Taylor BE, McClave SA, Martindale RG, Warren MM, Johnson DR, Braunschweig C, et al. Guidelines for the provision and assessment of nutrition support therapy in the adult critically ill patient. *Crit Care Med* 2016;44:390–438.
- [28] Mahadeva S, Malik A, Hilmi I, Qua C-S, Wong C-H, Goh K-L. Transnasal endoscopic placement of nasoenteric feeding tubes: outcomes and limitations in non-critically ill patients. *Nutr Clin Pract* 2008;23:176–81.
- [29] Dhaliwal R, Cahill N, Lemieux M, Heyland DK. The Canadian critical care nutrition guidelines in 2013. *Nutr Clin Pract* 2013;29:29–43.
- [30] Chowdhury AH, Murray K, Hoad CL, Costigan C, Marciani L, Macdonald IA, et al. Effects of bolus and continuous nasogastric feeding on gastric emptying, small bowel water content, superior mesenteric artery blood flow, and plasma hormone concentrations in healthy adults: a randomized crossover study. *Ann Surg* 2016;263:450–7.
- [31] Patel JJ, Rosenthal MD, Heyland DK. Intermittent versus continuous feeding in critically ill adults. *Curr Opin Clin Nutr Metab* 2018;21:116–20.
- [32] Compher C, Bingham AL, McCall M, Patel J, Rice TW, Braunschweig C, et al. Guidelines for the provision of nutrition support therapy in the adult critically ill patient: the American Society for Parenteral and Enteral Nutrition. *JPEN - J Parenter Enter Nutr* 2022;46:12–41.
- [33] Reignier J, Planteve G, Mira JP, Argaud L, Asfar P, Aissaoui N, et al. Low versus standard calorie and protein feeding in ventilated adults with shock: a randomised, controlled, multicentre, open-label, parallel-group trial (NUTRIREA-3). *Lancet Respir Med* 2023;11:602–12.
- [34] Pardo E, Lescot T, Preiser J-C, Massanet P, Pons A, Jaber S, et al. Association between early nutrition support and 28-day mortality in critically ill patients: the FRANS prospective nutrition cohort study. *Crit Care* 2023;27:7.
- [35] Reignier J, Boismamé-Helms J, Brisard L, Lascarrou JB, Ait Hssain A, Anguel N, et al. Enteral versus parenteral early nutrition in ventilated adults with shock: a randomised, controlled, multicentre, open-label, parallel-group study (NUTRIREA-2). *Lancet (London, England)* 2018;391:133–43.
- [36] Albrecht HC, Trawa M, Gretschesel S. Nonocclusive mesenteric ischemia associated with postoperative jejunal tube feeding: indicators for clinical management. *J Int Med Res* 2020;48:300060520929128.
- [37] Murthy TA, Plummer MP, Tan E, Chapman MJ, Chapple L-A. Higher versus lower enteral calorie delivery and gastrointestinal dysfunction in critical illness: a systematic review and meta-analysis. *Clin Nutr* 2022;41:2185–94.
- [38] Vijayaraghavan R, Maiwall R, Arora V, Choudhary A, Benjamin J, Aggarwal P, et al. Reversal of feed intolerance by prokinetics improves survival in critically ill cirrhosis patients. *Dig Dis Sci* 2022;67:4223–33.
- [39] Jenkins B, Calder PC, Marino LV. A systematic review of the definitions and prevalence of feeding intolerance in critically ill adults. *Clinical Nutrition ESPEN* 2022;49:92–102.
- [40] Bordejé ML, Montejo JC, Mateu ML, Solera M, Acosta JA, Juan M, et al. Intra-abdominal pressure as a marker of enteral nutrition intolerance in critically ill patients. The PIANE study. *Nutrients* 2019;11:2616.
- [41] Reintam Blaser A, Padar M, Mändul M, Elke G, Engel C, Fischer K, et al. Development of the Gastrointestinal Dysfunction Score (GIDS) for critically ill patients – a prospective multicenter observational study (iSOFA study). *Clin Nutr* 2021;40:4932–40.

- [42] Zusman O, Kagan I, Bendavid I, Theilla M, Cohen J, Singer P. Predictive equations versus measured energy expenditure by indirect calorimetry: a retrospective validation. *Clin Nutr* 2019;38:1206–10.
- [43] Frankenfield DC, Coleman A, Alam S, Cooney RN. Analysis of estimation methods for resting metabolic rate in critically ill adults. *J Parenter Enteral Nutr* 2008;33:27–36.
- [44] Tatucu-Babet OA, Ridley EJ, Tierney AC. Prevalence of underprescription or overprescription of energy needs in critically ill mechanically ventilated adults as determined by indirect calorimetry. *J Parenter Enteral Nutr* 2015;40:212–25.
- [45] Graf S, Pichard C, Genton L, Oshima T, Heidegger CP. Energy expenditure in mechanically ventilated patients: the weight of body weight. *Clin Nutr* 2017;36:224–8.
- [46] Singer P, Anbar R, Cohen J, Shapiro H, Shalita-Chesner M, Lev S, et al. The tight calorie control study (TICACOS): a prospective, randomized, controlled pilot study of nutritional support in critically ill patients. *Intensive Care Med* 2011;37:601–9.
- [47] Petros S, Horbach M, Seidel F, Weidhase L. Hypocaloric vs normocaloric nutrition in critically ill patients. *J Parenter Enteral Nutr* 2014;40:242–9.
- [48] Heidegger CP, Berger MM, Graf S, Zingg W, Darmon P, Costanza MC, et al. Optimisation of energy provision with supplemental parenteral nutrition in critically ill patients: a randomised controlled clinical trial. *Lancet* 2013;381:385–93.
- [49] Allingstrup MJ, Kondrup J, Wiis J, Claudius C, Pedersen UG, Hein-Rasmussen R, et al. Early goal-directed nutrition versus standard of care in adult intensive care patients: the single-centre, randomised, outcome assessor-blinded EAT-ICU trial. *Intensive Care Med* 2017;43:1637–47.
- [50] Stapel SN, de Grooth H-JS, Alimohamad H, Elbers PWG, Girbes ARJ, Weijls PJM, et al. Ventilator-derived carbon dioxide production to assess energy expenditure in critically ill patients: proof of concept. *Crit Care* 2015;19:370.
- [51] Oshima T, Graf S, Heidegger C-P, Genton L, Pugin J, Pichard C. Can calculation of energy expenditure based on CO(2) measurements replace indirect calorimetry? *Crit Care* 2017;21:13.
- [52] Berger MM, Pichard C. Understanding the Causes of Death in INTACT by Braunschweig et al. *J Parenter Enteral Nutr* 2015;39:144.
- [53] Heyland DK, Cahill N, Day AG. Optimal amount of calories for critically ill patients: depends on how you slice the cake. *Crit Care Med* 2011;39:2619–26.
- [54] Bellomo R, Cass A, Cole L, Finfer S, Gallagher M, Lee J, et al. Calorie intake and patient outcomes in severe acute kidney injury: findings from the Randomized Evaluation of Normal vs. Augmented Level of Replacement Therapy (RENAL) study trial. *Crit Care* 2014;18:R45 [R].
- [55] Weijls PJM, Looijaard WGP, Beishuizen A, Girbes ARJ, Oudemans-van Straaten HM. Early high protein intake is associated with low mortality and energy overfeeding with high mortality in non-septic mechanically ventilated critically ill patients. *Crit Care* 2014;18:701.
- [56] Zusman O, Theilla M, Cohen J, Kagan I, Bendavid I, Singer P. Resting energy expenditure, calorie and protein consumption in critically ill patients: a retrospective cohort study. *Crit Care* 2016;20:367.
- [57] Iapichino G, Radrizzani D, Armani S, Noto A, Spanu P, Mistràletti G. Metabolic trend of critically ill patients: energy balance and substrate disposal. *Minerva Anestesiol* 2006;72:533–41.
- [58] Dvir D, Cohen J, Singer P. Computerized energy balance and complications in critically ill patients: an observational study. *Clin Nutr* 2006;25:37–44.
- [59] Villet S, Chiolerio RL, Bollmann MD, Revelly J-P, Cayeux Rn M-C, Delarue J, et al. Negative impact of hypocaloric feeding and energy balance on clinical outcome in ICU patients. *Clin Nutr* 2005;24:502–9.
- [60] Heyland DK, Patel J, Compher C, Rice TW, Bear DE, Lee ZY, et al. The effect of higher protein dosing in critically ill patients with high nutritional risk (EFFORT Protein): an international, multicentre, pragmatic, registry-based randomised trial. *Lancet* (London, England) 2023;401:568–76.
- [61] Lin J, Chen W, Ye X, Lv C, Liu Y, Jiang X, et al. Trajectories of protein intake and 28-day mortality in critically ill patients: a secondary analysis of a cluster-randomized controlled trial. *Clin Nutr* 2022;41:1644–50.
- [62] Koekkoek WAC, van Setten CH, Olthoff LE, Kars JCN, van Zanten ARH. Timing of PROTein INtake and clinical outcomes of adult critically ill patients on prolonged mechanical Ventilation: the PROTINVENT retrospective study. *Clin Nutr* 2019;38:883–90.
- [63] Gamrin-Gripenberg L, Sundström-Rehal M, Olsson D, Grip J, Wernerman J, Rooyackers O. An attenuated rate of leg muscle protein depletion and leg free amino acid efflux over time is seen in ICU long-stayers. *Crit Care* 2018;22:13.
- [64] Deutz NEP, Singer P, Wierchowska-McNew RA, Viana MV, Ben-David IA, Pantet O, et al. Comprehensive metabolic amino acid flux analysis in critically ill patients. *Clin Nutr* 2021;40:2876–97.
- [65] Burtin C, Clerckx B, Robbeets C, Ferdinande P, Langer D, Troosters T, et al. Early exercise in critically ill patients enhances short-term functional recovery. *Crit Care Med* 2009;37:2499–505.
- [66] Schaller SJ, Anstey M, Blobner M, Edrich T, Grabitz SD, Gradwohl-Matis I, et al. Early, goal-directed mobilisation in the surgical intensive care unit: a randomised controlled trial. *Lancet* 2016;388:1377–88.
- [67] Bier DM, Brosnan JT, Flatt JP, Hanson RW, Heird W, Hellerstein MK, et al. Report of the IDECG Working Group on lower and upper limits of carbohydrate and fat intake. *Eur J Clin Nutr* 1999;53:s177–8.
- [68] Thorell A, Rooyackers O, Myrenfors P, Soop M, Nygren J, Ljungqvist OH. Intensive insulin treatment in critically ill trauma patients normalizes glucose by reducing endogenous glucose production. *J Clin Endocrinol Metabol* 2004;89:5382–6.
- [69] Tappy L, Schwarz J-M, Schneiter P, Cayeux C, Revelly J-P, Fagerquist CK, et al. Effects of isoenergetic glucose-based or lipid-based parenteral nutrition on glucose metabolism, de novo lipogenesis, and respiratory gas exchanges in critically ill patients. *Crit Care Med* 1998;26:860–7.
- [70] Han Y-Y, Lai S-R, Partridge JS, Wang MY, Sulo S, Tsao F-W, et al. The clinical and economic impact of the use of diabetes-specific enteral formula on ICU patients with type 2 diabetes. *Clin Nutr* 2017;36:1567–72.
- [71] Mesejo A, Montejo-González JC, Vaquerizo-Alonso C, Lobo-Tamer G, Zabarte-Martínez M, Herrero-Meseguer JL, et al. Diabetes-specific enteral nutrition formula in hyperglycemic, mechanically ventilated, critically ill patients: a prospective, open-label, blind-randomized, multicenter study. *Crit Care* 2015;19:390.
- [72] Green P, Theilla M, Singer P. Lipid metabolism in critical illness. *Curr Opin Clin Nutr Metab Care* 2016;19:111–5.
- [73] Elwyn DH. Protein and energy requirements: effect of clinical state. *Clin Nutr* 1993;12:S44–51.
- [74] Berger MM, Reintam-Blaser A, Calder PC, Casaer M, Hiesmayr MJ, Mayer K, et al. Monitoring nutrition in the ICU. *Clin Nutr* 2019;38:584–93.
- [75] Bousie E, van Blokland D, Lammers HJW, van Zanten ARH. Relevance of non-nutritional calories in mechanically ventilated critically ill patients. *Eur J Clin Nutr* 2016;70:1443–50.
- [76] Calder PC, Adolph M, Deutz NE, Grau T, Innes JK, Klek S, et al. Lipids in the intensive care unit: recommendations from the ESPEN expert group. *Clin Nutr* 2018;37:1–18.
- [77] Fürst P, Albers S, Stehle P. Availability of glutamine supplied intravenously as alanylglutamine. *Metabolism* 1989;38:67–72.
- [78] Rousseau A-F, Losser M-R, Ichaï C, Berger MM. ESPEN endorsed recommendations: nutritional therapy in major burns. *Clin Nutr* 2013;32:497–502.
- [79] Heyland DK, Wibbenmeyer L, Pollack JA, Friedman B, Turgeon AF, Eshraghi N, et al. A randomized trial of enteral glutamine for treatment of burn injuries. *N Engl J Med* 2022;387:1001–10.
- [80] Gonzalez MR, Fleuchot B, Lauciello L, Jafari P, Applegate LA, Raffoul W, et al. Effect of human burn wound exudate on *Pseudomonas aeruginosa* virulence. *mSphere* 2016;1:00111–5.
- [81] Azman M, Mohd Yunus MR, Sulaiman S, Syed Omar SN. Enteral glutamine supplementation in surgical patients with head and neck malignancy: a randomized controlled trial. *Head Neck* 2015;37:1799–807.
- [82] Rodas PC, Rooyackers O, Hebert C, Å Norberg, Wernerman J. Glutamine and glutathione at ICU admission in relation to outcome. *Clin Sci (Lond)* 2012;122:591–7.
- [83] van Zanten ARH, Sztark F, Kaisers UX, Zielmann S, Felbinger TW, Sablotzki AR, et al. High-protein enteral nutrition enriched with immune-modulating nutrients vs standard high-protein enteral nutrition and nosocomial infections in the ICU. *JAMA* 2014;312:514.
- [84] Pasin L, Landoni G, Zangrillo A. Glutamine and antioxidants in critically ill patients. *N Engl J Med* 2013;369:482–4.
- [85] Heyland DK, Elke G, Cook D, Berger MM, Wischmeyer PE, Albert M, et al. Glutamine and antioxidants in the critically ill patient: a post hoc analysis of a large-scale randomized trial. *JPEN - J Parenter Enter Nutr* 2015;39:401–9.
- [86] Stehle P, Ellger B, Kojic D, Feuersenger A, Schneid C, Stover J, et al. Glutamine dipeptide-supplemented parenteral nutrition improves the clinical outcomes of critically ill patients: a systematic evaluation of randomised controlled trials. *Clinical Nutrition ESPEN* 2017;17:75–85.
- [87] Zhu D, Zhang Y, Li S, Gan L, Feng H, Nie W. Enteral omega-3 fatty acid supplementation in adult patients with acute respiratory distress syndrome: a systematic review of randomized controlled trials with meta-analysis and trial sequential analysis. *Intensive Care Med* 2014;40:504–12.
- [88] Glenn JOH, Wischmeyer PE. Enteral fish oil in critical illness. *Curr Opin Clin Nutr Metab Care* 2014;17:116–23.
- [89] Kagan I, Cohen J, Stein M, Bendavid I, Pinsker D, Silva V, et al. Preemptive enteral nutrition enriched with eicosapentaenoic acid, gamma-linolenic acid and antioxidants in severe multiple trauma: a prospective, randomized, double-blind study. *Intensive Care Med* 2015;41:460–9.
- [90] Hofman Z, Swinkels S, van Zanten ARH. Glutamine, fish oil and antioxidants in critical illness: MetaPlus trial post hoc safety analysis. *Ann Intensive Care* 2016;6:119.
- [91] Calder PC, Waitzberg DL, Klek S, Martindale RG. Lipids in parenteral nutrition: biological aspects. *JPEN - J Parenter Enter Nutr* 2020;44(Suppl 1):S21. s7.
- [92] Singer P, Calder PC. The role of omega-3 polyunsaturated fatty acids in the intensive care unit. *Curr Opin Clin Nutr Metab Care* 2023;26:129–37.
- [93] Pradelli L, Mayer K, Klek S, Omar Alsaleh AJ, Clark RAC, Rosenthal MD, et al. ω-3 fatty-acid enriched parenteral nutrition in hospitalized patients: systematic review with meta-analysis and trial sequential analysis. *JPEN - J Parenter Enter Nutr* 2020;44:44–57.

- [94] Wang C, Han D, Feng X, Wu J. Omega-3 fatty acid supplementation is associated with favorable outcomes in patients with sepsis: an updated meta-analysis. *J Int Med Res* 2020;48:300060520953684.
- [95] Wang H, Su S, Wang C, Hu J, Dan W, Peng X. Effects of fish oil-containing nutrition supplementation in adult sepsis patients: a systematic review and meta-analysis. *Burns & trauma* 2022;10:tkac012.
- [96] Pradelli L, Klek S, Mayer K, Omar Alsaleh AJ, Rosenthal MD, Heller AR, et al. Omega-3 fatty acid-containing parenteral nutrition in ICU patients: systematic review with meta-analysis and cost-effectiveness analysis. *Crit Care* 2020;24:634.
- [97] Hoeger J, Simon T-P, Beeker T, Marx G, Haase H, Schuerholz T. Persistent low serum zinc is associated with recurrent sepsis in critically ill patients - a pilot study. *PLoS One* 2017;12:e0176069 [e].
- [98] Ben-Hamouda N, Charrière M, Voirol P, Berger MM. Massive copper and selenium losses cause life-threatening deficiencies during prolonged continuous renal replacement. *Nutrition* 2017;34:71–5.
- [99] Manzanara W, Lemieux M, Elke G, Langlois PL, Bloos F, Heyland DK. High-dose intravenous selenium does not improve clinical outcomes in the critically ill: a systematic review and meta-analysis. *Crit Care* 2016;20:356.
- [100] Han JH, Ginde AA, Brown SM, Baughman A, Collar EM, Ely EW, et al. Effect of early high-dose vitamin D3 repletion on cognitive outcomes in critically ill adults. *Chest* 2021;160:909–18.
- [101] Ginde AA, Brower RG, Caterino JM, Finck L, Banner-Goodspeed VM, Grisson CK, et al. Blood Institute, Petal Clinical Trials Network. Early high-dose vitamin D(3) for critically ill, vitamin D-deficient patients. *N Engl J Med* 2019;381:2529–40.
- [102] Brown J, Robertson C, Sevilla L, Garza J, Rashid H, Benitez AC, et al. A systematic review and meta-analysis on possible role of vitamin C in sepsis. *Cureus* 2022;14:e32886 [e].
- [103] Chen C-Y, Chiu C-T, Lee H-S, Lai C-C. The impact of vitamin C-containing treatment on the mortality of patients with sepsis: a systematic review and meta-analysis of randomized controlled trials. *Journal of Infection and Public Health* 2022;15:1514–20.
- [104] Lamontagne F, Masse M-H, Menard J, Sprague S, Pinto R, Heyland DK, et al. Intravenous vitamin C in adults with sepsis in the intensive care unit. *N Engl J Med* 2022;386:2387–98.
- [105] Langlois PL, Manzanara W, Adhikari NKJ, Lamontagne F, Stoppe C, Hill A, et al. Vitamin C administration to the critically ill: a systematic review and meta-analysis. *J Parenter Enteral Nutr* 2018;43:335–46.
- [106] Lee Z-Y, Ortiz-Reyes L, Lew CCH, Hasan MS, Ke L, Patel JJ, et al. Intravenous vitamin C monotherapy in critically ill patients: a systematic review and meta-analysis of randomized controlled trials with trial sequential analysis. *Ann Intensive Care* 2023;13:14.
- [107] Martimbianco ALC, Pacheco RL, Bagattini ÂM, de Fátima Carreira Moreira Padovez R, Azevedo LCP, Riera R. Vitamin C-based regimens for sepsis and septic shock: systematic review and meta-analysis of randomized clinical trials. *J Crit Care* 2022;71:154099.
- [108] Wen C, Li Y, Hu Q, Liu H, Xu X, Lü M. IV vitamin C in sepsis: a latest systematic review and meta-analysis. *Int J Clin Pract* 2023;2023:6733465.
- [109] Lu D, Mao W. Hydrocortisone combined with vitamin C and thiamine in the treatment of sepsis/septic shock: a systematic review with meta-analysis and trial sequential analysis. *Clin Invest Med* 2023;46:E36–49.
- [110] Zajic P, Amrein K. Vitamin D deficiency in the ICU: a systematic review. *Minerva Endocrinol* 2014;39:275–87.
- [111] Reeves A, White H, Sosnowski K, Tran K, Jones M, Palmer M. Energy and protein intakes of hospitalised patients with acute respiratory failure receiving non-invasive ventilation. *Clin Nutr* 2014;33:1068–73.
- [112] Tsai M-H, Ku S-C, Wang T-G, Hsiao T-Y, Lee J-J, Chan D-C, et al. Swallowing dysfunction following endotracheal intubation: age matters. *Medicine (Baltimore)* 2016;95:e3871 [e].
- [113] Macht M, Wimbish T, Clark BJ, Benson AB, Burnham EL, Williams A, et al. Postextubation dysphagia is persistent and associated with poor outcomes in survivors of critical illness. *Crit Care* 2011;15:R231 [R].
- [114] Macht M, White SD, Moss M. Swallowing dysfunction after critical illness. *Chest* 2014;146:1681–9.
- [115] Pryor L, Ward E, Cornwell P, O'Connor S, Chapman M. Patterns of return to oral intake and decannulation post-tracheostomy across clinical populations in an acute inpatient setting. *Int J Lang Commun Disord* 2016;51:556–67.
- [116] Fujita T, Daiko H, Nishimura M. Early enteral nutrition reduces the rate of life-threatening complications after thoracic esophagectomy in patients with esophageal cancer. *Eur Surg Res* 2012;48:79–84.
- [117] Melis M. Bowel necrosis associated with early jejunal tube feeding. *Arch Surg* 2006;141:701.
- [118] Sarap AN, Sarap MD, Childers J. Small bowel necrosis in association with jejunal tube feeding. *J Am Acad Physician Assistants* 2010;23:28.
- [119] Pupelis G, Selga G, Austrums E, Kaminski A. Jejunal feeding, even when instituted late, improves outcomes in patients with severe pancreatitis and peritonitis. *Nutrition* 2001;17:91–4.
- [120] Malhotra A, Mathur AK, Gupta S. Early enteral nutrition after surgical treatment of gut perforations: a prospective randomised study. *J Postgrad Med* 2004;50:102–6.
- [121] Kaur N, Gupta MK, Minocha VR. Early enteral feeding by nasoenteric tubes in patients with perforation peritonitis. *World J Surg* 2005;29:1023–7.
- [122] Picot D, Layec S, Dussaux L, Trivin F, Thibault R. Chyme reinfusion in patients with intestinal failure due to temporary double enterostomy: a 15-year prospective cohort in a referral centre. *Clin Nutr* 2017;36:593–600.
- [123] Goiburu ME, Goiburu MM, Bianco H, Díaz JR, Alderete F, Palacios MC, et al. The impact of malnutrition on morbidity, mortality and length of hospital stay in trauma patients. *Nutr Hosp* 2006;21:604–10.
- [124] Kompan L, Vidmar G, Spindler-Vesel A, Pečar J. Is early enteral nutrition a risk factor for gastric intolerance and pneumonia? *Clin Nutr* 2004;23:527–32.
- [125] Fan M-c, Wang Q-l, Fang W, Jiang Y-x, Li L-d, Sun P, et al. Early enteral combined with parenteral nutrition treatment for severe traumatic brain injury: effects on immune function, nutritional status and outcomes. *Chin Med Sci J* 2016;31:213–20.
- [126] Pan J, Shaffer R, Sinno Z, Tyler M, Ghosh J. The obesity paradox in ICU patients. In: 2017 39th annual international conference of the IEEE engineering in medicine and biology society (EMBC). IEEE; 2017.
- [127] Dickerson RN, Patel JJ, McClain CJ. Protein and calorie requirements associated with the presence of obesity. *Nutr Clin Pract* 2017;32:865–93S.
- [128] Elke G, Wang M, Weiler N, Day AG, Heyland DK. Close to recommended caloric and protein intake by enteral nutrition is associated with better clinical outcome of critically ill septic patients: secondary analysis of a large international nutrition database. *Crit Care* 2014;18:R29 [R].
- [129] Chuntrasakul C, Siltharm S, Chinswangwatanakul V, Pongprasobchai T, Chockvivatanavanit S, Bunnak A. Early nutritional support in severe traumatic patients. *Journal of the Medical Association of Thailand = Chotmaihet thangphaet* 1996;79:21–6.
- [130] Pupelis EAJG. Randomised trial of safety and efficacy of postoperative enteral feeding in patients with severe pancreatitis: preliminary report. *Eur J Surg* 2000;166:383–7.
- [131] Manc EE, Muzevich KM. Tolerability and safety of enteral nutrition in critically ill patients receiving intravenous vasopressor therapy. *J Parenter Enteral Nutr* 2012;37:641–51.
- [132] Khalid I, Doshi P, DiGiovine B. Early enteral nutrition and outcomes of critically ill patients treated with vasopressors and mechanical ventilation. *Am J Crit Care* 2010;19:261–8.
- [133] Wang Z, Heshka S, Gallagher D, Boozer CN, Kotler DP, Heymsfield SB. Resting energy expenditure-fat-free mass relationship: new insights provided by body composition modeling. *Am J Physiol Endocrinol Metabol* 2000;279:E539–45.
- [134] Dickerson RN. Hypocaloric, high-protein nutrition therapy for critically ill patients with obesity. *Nutr Clin Pract* 2014;29:786–91.
- [135] Viana MV, Moraes RB, Fabbri AR, Santos MF, Torman VBL, Vieira SR, et al. Contrasting effects of preexisting hyperglycemia and higher body size on hospital mortality in critically ill patients: a prospective cohort study. *BMC Endocr Disord* 2014;14:50.
- [136] Egi M, Bellomo R, Stachowski E, French CJ, Hart G. Variability of blood glucose concentration and short-term mortality in critically ill patients. *Anesthesiology* 2006;105:244–52.
- [137] Hermanides J, Vriesendorp TM, Bosman RJ, Zandstra DF, Hoekstra JB, DeVries JH. Glucose variability is associated with intensive care unit mortality. *Crit Care Med* 2010;38:838–42.
- [138] Egi M, Krinsley JS, Maurer P, Amin DN, Kanazawa T, Ghandi S, et al. Pre-morbid glycemic control modifies the interaction between acute hypoglycemia and mortality. *Intensive Care Med* 2016;42:562–71.
- [139] Yatabe T, Inoue S, Sakaguchi M, Egi M. The optimal target for acute glycemic control in critically ill patients: a network meta-analysis. *Intensive Care Med* 2016;43:16–28.
- [140] Ichai C, Preiser J-C, Société Française d'A-R, Société de Réanimation de Langue F, Experts g. International recommendations for glucose control in adult non diabetic critically ill patients. *Crit Care* 2010;14:R166 [R].
- [141] Schultz MJ, Harmsen RE, Spronk PE. Clinical review: strict or loose glycemic control in critically ill patients-implementing best available evidence from randomized controlled trials. *Crit Care* 2010;14:223.
- [142] Ridley EJ, Davies AR. Practicalities of nutrition support in the intensive care unit: the usefulness of gastric residual volume and prokinetic agents with enteral nutrition. *Nutrition* 2011;27:509–12.
- [143] Reigner J. Effect of not monitoring residual gastric volume on risk of ventilator-associated pneumonia in adults receiving mechanical ventilation and early enteral feeding. *JAMA* 2013;309:249.
- [144] Rhodes A, Evans LE, Alhazzani W, Levy MM, Antonelli M, Ferrer R, et al. Surviving sepsis campaign: international guidelines for management of sepsis and septic shock: 2016. *Intensive Care Med* 2017;43:304–77.
- [145] Mehanna HM, Moledina J, Travis J. Refeeding syndrome: what it is, and how to prevent and treat it. *BMJ* 2008;336:1495–8.
- [146] Rio A, Whelan K, Goff L, Reidlinger DP, Smeeton N. Occurrence of refeeding syndrome in adults started on artificial nutrition support: prospective cohort study. *BMJ Open* 2013;3:e002173.
- [147] Doig GS, Simpson F, Heighes PT, Bellomo R, Cheshire D, Caterson ID, et al. Restricted versus continued standard caloric intake during the management of refeeding syndrome in critically ill adults: a randomised, parallel-group, multicentre, single-blind controlled trial. *Lancet Respir Med* 2015;3:943–52.
- [148] Koekkoek KWAC, van Zanten ARH. Nutrition in the ICU. *Curr Opin Anaesthesiol* 2018;31:136–43.
- [149] Olthof LE, Koekkoek WACK, van Setten C, Kars JCN, van Blokland D, van Zanten ARH. Impact of caloric intake in critically ill patients with, and without, refeeding syndrome: a retrospective study. *Clin Nutr* 2018;37:1609–17.