

As a library, NLM provides access to scientific literature. Inclusion in an NLM database does not imply endorsement of, or agreement with, the contents by NLM or the National Institutes of Health.

Learn more: [PMC Disclaimer](#) | [PMC Copyright Notice](#)



Nutrients. 2024 Mar 25;16(7):941. doi: [10.3390/nu16070941](https://doi.org/10.3390/nu16070941)

Comparative Efficacy of Different Protein Supplements on Muscle Mass, Strength, and Physical Indices of Sarcopenia among Community-Dwelling, Hospitalized or Institutionalized Older Adults Undergoing Resistance Training: A Network Meta-Analysis of Randomized Controlled Trials

[Chun-De Liao](#)^{1,2,†}, [Shih-Wei Huang](#)^{2,3,†}, [Hung-Chou Chen](#)^{2,3}, [Mao-Hua Huang](#)⁴, [Tsan-Hon Liou](#)^{2,3}, [Che-Li Lin](#)^{5,6,*}

Editors: Mario Barbagallo, Luana Toniolo, Emiliana Giacomello

[Author information](#) [Article notes](#) [Copyright and License information](#)

PMCID: PMC11013298 PMID: [38612975](#)

Abstract

Aging-related sarcopenia exerts harmful impacts on muscle mass, strength, and physical mobility. Protein supplementation has been demonstrated to augment efficacy of resistance training (RT) in elderly. This study compared the relative effects of different protein supplements on muscle mass, strength, and mobility outcomes in middle-aged and older individuals undergoing RT. A comprehensive search of online databases was performed to identify randomized controlled trials (RCTs) examining the efficacy of protein supplement plus RT in untrained community-dwelling adults, hospitalized, or institutionalized residents who suffered acute or chronic health conditions. Network meta-analysis (NMA) was performed using a frequentist method for all analyses. Treatment effects for main outcomes were expressed as standard mean difference (SMD) with 95% confidence interval (CI). We used the surface-under-the cumulative-ranking (SUCRA) scores to rank probabilities of effect estimation among all identified treatments. Meta-regression analyses were performed to identify any relevant moderator of the treatment efficacy and results were expressed as β with 95% credible interval (CrI). We finally included 78 RCTs (5272 participants) for analyses. Among the six protein sources identified in this NMA, namely whey, milk, casein, meat, soy, and peanut, whey supplement yielded the most effective treatments augmenting efficacy of RT on muscle mass (SMD = 1.29, 95% CI: 0.96, 1.62; SUCRA = 0.86), handgrip strength (SMD = 1.46, 95% CI: 0.92, 2.00; SUCRA = 0.85), and walking speed (SMD = 0.73, 95% CI: 0.39, 1.07; SUCRA = 0.84). Participant's health condition, sex, and supplementation dose were significant factors moderating the treatment efficacy on muscle mass (β = 0.74; 95% CrI: 0.22, 1.25), handgrip strength (β = −1.72; 95% CrI:

−2.68, −0.77), and leg strength ($\beta = 0.76$; 95% CrI: 0.06, 1.47), respectively. Our findings suggest whey protein yields the optimal supplements to counter sarcopenia in older individuals undergoing RT.

Keywords: sarcopenia, protein supplement, resistance exercise, muscle mass, strength, mobility

1. Introduction

Sarcopenia, an age-driven skeletal muscle condition, has been recognized as a muscle disease characterized by progressive attenuation of muscle mass [1,2]. The loss of muscle mass can further impact muscle strength and its function performance which in turn results in high risks of physical difficulty and disability in elderly [3,4,5,6]. Senile primary sarcopenia commonly occurs with advancing age which is accompanied with a number of chronic comorbidities or long-term health conditions such as cardiovascular diseases [7], arthritis [8,9,10,11], diabetes mellitus [12], and stroke [13]; accordingly, sarcopenia has become epidemic in the ageing population. Depending on the various diagnostic criteria, 5–13% of people who aged older than 60 years are identified as having sarcopenia, with an increasing prevalence up to 50% in elderly aged older than 80 years [14,15]. In worldwide, an overall estimate of prevalence of sarcopenia is reported to be around 10% in community-dwelling older individuals, 31–51% in long-term care institutions and 24% in the acute hospital-care population [16,17]. Because of the growing numbers on prevalence and the accompanied adverse effects of sarcopenia [18,19,20,21,22], such disease is responsible for considerable burden in healthcare system [23,24]. Therefore, it is imperative to investigate effective treatment strategies developed to preserve or restore muscle mass as early as possible for elderly people.

The primary sarcopenia is defined based on handgrip strength, lean mass, and mobility due to progressive skeletal muscle wasting and related physical decline with aging. The major underlying mechanism driving the disease development and progression of sarcopenia has been addressed to the blunted muscle protein synthetic response (i.e., myogenesis) and negative muscle net balance in elder people [25,26,27]. Additionally, sarcopenia is commonly recognized as a multi-etiological disease associated with a multiplicity of physical, psychological, and social factors in community-dwelling older adults, among which physical inactivity as well as malnutrition are determinant factors that have great impacts on sarcopenia [28]. Therefore, the most effective managements for sarcopenia rely on the adoption of healthy lifestyle behaviors, including adherence to regular physical activity and high-quality diets [29,30,31,32]. Under such scenario, the optimal treatment strategy of sarcopenia comprises multidisciplinary interventions including exercise and nutrition treatment [33,34,35]. With respect to physical activity intervention, resistance training (RT) has been identified as a successful exercise therapy which have promising effects on counteracting sarcopenia in terms of increased whole-body lean mass and muscular strength in elderly population [36,37,38,39]. Aside from regular physical activity, nutritional interventions using nutrient supplements are targeted to prevent or postpone the onset of such geriatric conditions derived from sarcopenia [32,40,41]. Especially, a high-protein diet or protein supplementation is believed to have surplus benefits in augmenting efficacy of RT for healthy older people [42,43,44,45], as well as those who have high risks of sarcopenia [41,43,44,45,46,47,48,49,50,51]. Sufficient protein intake plays a key role to stimulate maximal rate of muscle protein synthesis leading to anabolism and net new muscle protein accretion during RT [52,53]. Such resultant net of muscle protein is in turn accumulated over the training period to yield muscle mass gain or morphological hypertrophy.

The postprandial muscle protein synthetic response appears to be modulated by the amount, source, and type of protein consumed [54]. Numerous protein nutrients are available for individuals to intake and meet their daily protein needs, including animal-derived (e.g., whey, casein, milk, or beef etc.) and plant-derived (e.g., soy, pea, or rice etc.) dietary protein sources, and most of which have been shown to effectively stimulate post-exercise myofibrillar protein synthesis rates [55,56,57,58]. Due to that rate of digestibility and amount of leucine content are varied among different protein supplements, the muscle protein synthesis rates in response to the ingestion of such protein supplements may differ at rest and following RT. Empirical support for this inference have been observed [59,60,61]. Therefore, in terms of muscle protein synthesis rate, the protein source and type deem an important factor determining the acute treatment efficacy of such intervention of nutrient supplementation along with RT. However, it remains unclear whether the variety type or source of protein supplements obtain different long-term outcome of muscle mass and its function after RT intervention. Identifying relative treatment efficacy among different protein supplements assists the clinical practitioners establishing optimal and efficient intervention strategies.

Previous systemic review and meta-analysis studies have compared the effects of different protein sources during exercise training [44,62,63,64,65,66,67], among which the compared protein supplements included whey [44,62,63,64,65,67], milk [44,63,65,67], casein [44,62,65,67], beef [63,64], soy [62,63,65,67], and cereal protein [63,65]. However, all such protein supplements were not yet fully compared in single meta-analysis study and the most effective treatment option remains unknown. In addition, most of the previous systemic reviews enrolled both young and older adults, only few targeted the elderly population alone [44,65,67]. Based on the increasing number of sarcopenic individual among elderly population, there is an urgent need to identify the most effective and efficient treatment strategy for such population restoring their muscle strength and function. With respect to the above considerations, the purpose of this study was to (1) compare the relative efficacy of different type of protein supplements combined with RT on muscle mass, strength, and physical mobility by using network meta-analysis (NMA); (2) identify if there is any relevant moderator affecting the relative treatment efficacy; and (3) determine the optimal treatment by ranking the probability of each supplement type for community-dwelling or institutionalized older adults undergoing RT.

2. Materials and Methods

2.1. Study Protocol

This NMA adapted the guidelines of the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) [68] and the extension statement for reporting of systematic reviews incorporating NMA [69,70]. The protocol of the present NMA was registered at PROSPERO (registration number: CRD42023472942).

2.2. Search Strategy

Electronic databases including PubMed, the Cochrane Library Database, EMBASE, the ClinicalKey, the China Knowledge Resource Integrated Database, the Physiotherapy Evidence Database, and Google Scholar were searched from inception to 31 July 2023 to identify eligible randomized controlled trials (RCTs) investigating the efficacy of protein supplementation (PS), resistance training (RT), or combined treatment for older people.

We also performed manual search for RCTs cited in articles retrieved from the databases. There was no limitation for the publication year or language.

The following search terms were based on PICOS (Participants, Intervention, Control, Outcomes, Study design) criteria: (1) Participants' conditions: ("older adult" OR "elder individual") AND ("untrained" OR "sedentary" OR "frailty" OR "sarcopenia" OR "nursing home" OR "institutionalization" OR "hospitalization"). (2) Intervention: ("resistance exercise training" OR "strengthening exercise") AND ("protein supplementation" OR "high-protein diet"); (3) Comparisons: placebo supplementation with or without RT, RT alone, usual medication or regular care (RC) without any PS or RT; (4) Outcomes: lean mass OR strength OR mobility. The details for main outcome measures were described in the following [Section 2.4](#); (5) Study design: a parallel or crossover RCT with two or multiple study arms. The detailed search formulas for each database are presented in [Supplementary Table S1](#).

2.3. Criteria for Selecting Studies

Trials were included if they met the following criteria: (1) the study design was an RCT (2) the study enrolled institutionalized residents or community-dwelling elders who aged ≥ 50 years, were untrained, sedentary, frail, or identified as having sarcopenia. Participants were excluded if they had considerable conditions in hematological, hepatic, or renal functions; (3) the treatment groups received PS in combination with RT. The PS involved different type of protein supplements derived from animal or plant sources; (4) the comparator group received an active treatment including comparative PS alone, placebo supplementation, RT with or without placebo supplementation, or regular care (RC) which was not related to PS or RT; and (5) the study reported main outcome measures defined in the following [Section 2.4](#).

The exclusion criteria were as follows: (1) a vitro or vivo study; (2) a case report or case series study; and (3) a prospective study without randomised allocation.

Study search, screen, and selection for relevant studies were independently performed by two authors (CDL and SWH). Another team members (THL and CLL) were consulted for any disagreement between the authors until consensus reached.

2.4. Main Outcomes

The main outcomes in this NMA included measures of muscle mass, muscle strength, and physical mobility, all of which are diagnostic indicators of sarcopenia recommended by the Asian Working Group for Sarcopenia [\[1\]](#) and the European Working Group on Sarcopenia in Older People [\[2\]](#). The measures assessing skeletal muscle mass and volume included, but were not limited to, whole body lean mass, fat-free mass, appendicular lean mass, muscle thickness, muscle cross-sectional area, and circumference of arms or legs. Regarding muscle strength measures, handgrip strength was extracted as priori [\[2\]](#). The strength measures for lower extremity were extracted based on the following sequence of preference: Peak torque and maximum voluntary isometric contraction of quadriceps hamstring muscles, and hip abductors. Physical mobility was assessed using walking speed, chair rising ability, timed up-and-go (TUG) task, or the Short Physical Performance Battery (SPPB) score.

2.5. Data Extraction

We established a worksheet to extract the following data from each included study: (1) characteristics of study arm according to the intervention nature; (2) characteristics of the participants including age, sex distribution, and body mass index (BMI); (3) characteristics of the PS and RT protocols; (4) follow-up time points; and (5) main outcome measures. The relevant data from included trials were initially extracted by the author (CDL), and another author (SWH) validated and confirmed the extracted data. In cases of inconclusiveness, the author (CLL) was consulted to discuss the disagreements in a consensus meeting.

Subgroup analysis was performed based on the follow-up time frame which was defined as short-term (≤ 3 months), medium-term (> 3 months and ≤ 6 months), and long-term (> 6 months) for each outcome; when multiple time points were conducted within the same time frame, the longest one was selected for analysis (e.g., if the follow-up time points for muscle mass measurement were 6 and 12 weeks, the data measured at the 3-month time point were used as the short-term results). We also examined the compliance as well as side effects responding to PS and RT interventions among the included studies.

2.6. Risks of Bias in Individual Study and across Studies

Two reviewers (CDL and SWH) independently assessed the risk of bias of the included studies using the Cochrane Collaboration's risk-of-bias tool for quality assessment [71,72]. Any discrepancy between reviewers' assessments was resolved by a consensus meeting and, if necessary, discussed with a third reviewer (THL) until consensus was reached. The following six domains seven judgement items related to biased estimates of treatment effects were assessed [72]: (1) the selection bias determined by two judgements regarding random sequence generation and allocation concealment; (2) the performance bias judged by blinding of participants; (3) the detection bias associated with blind of outcome assessors; (4) the attrition bias corresponded with incomplete outcome data; (5) the reporting bias raised from selective reporting; and (6) other bias considering other sources of potential bias such as lack of information on sponsors, funding sources, or author conflict of interest disclosures which raise agenda bias. Each item was judged as low-risk, unclear-risk (specific details or description were not reported) or high-risk (not fulfilling the criteria) according to criteria specified by Cochrane. Based on the judged scores of the six domains of bias, an overall bias was identified as low, unclear, or high risk for each study [72].

To assess publication bias for each outcome, a funnel plot was employed to visually inspect any asymmetry among all effect sizes of treatment arms. In addition, we performed the Begg–Mazumdar rank correlation test to identify any significant publication bias across all included RCTs in each outcome [73].

2.7. Data Synthesis and Analysis

Treatment effects were expressed as standard mean differences (SMDs) alongside 95% confidence intervals (CIs) of each main outcome across the included RCTs. The mean value of change scores from baseline and its standard deviation (SD) were directly extracted from each study arm of the included RCTs. The missing SD of change

score was estimated using the SD of baseline (SD_{base}) and posttest (SD_{post}) measured data by the following calculation:

$$SD_{\text{change}} = \sqrt{\left[\left(SD_{\text{base}} \right)^2 + \left(SD_{\text{post}} \right)^2 - 2 \times \text{Corr}_{\text{base-post}} \times SD_{\text{base}} \times SD_{\text{post}} \right]}$$

Based on the recommended method [74], the within-participant correlation coefficient ($\text{Corr}_{\text{base-post}}$) represents the mean of the available correlations from the included studies in which the SD change was available. Accordingly, we used the resulted correlation coefficient of 0.9 for muscle mass measures and 0.7 for strength and mobility measures. Wherever the SD was not presented, it was estimated using *p*-values or 95% CIs. If data were reported as the median, range, or interquartile range, we followed the Wan's methods to estimate the mean value [75]; and accordingly, the SD was estimated by dividing the interquartile range by 1.35 [74]. The magnitude of the SMD was interpreted based on Cohen's criteria of effect size (*d*) and was categorized in accordance with the criteria [76]: trivial ($d < 0.10$), small ($0.10 \leq d < 0.25$), medium ($0.25 \leq d < 0.40$), and large ($d \geq 0.40$).

A random-effects NMA model was performed using a frequentist method to estimate effects from direct and indirect comparisons among all treatment arms. Heterogeneity as well as global consistency were assessed using the I^2 statistic alongside τ^2 values to estimate variance across the comparisons. The node-splitting method was used to assess consistency between direct and indirect comparisons and results were resented in a forest plot [77]. In addition, a full design-by-treatment interaction model was established to test total inconsistency across all treatment arms [78]. Ranking probabilities of effect estimation among all treatment arms per outcome were performed and expressed as the surface under the cumulative ranking (SUCRA) score [79].

We established the network meta-regression (NMR) models to identify any relevant moderators affecting the relative efficacy among treatment options. For each main outcome, the univariate NMR model was using an individual moderator as a covariate performed accordingly [80]. Substantial moderators were identified on the basis of (1) participant characteristics including age, BMI, female sex distribution of study sample (i.e., percentage proportion of women of total number of participants), world region of study application, and health status (i.e., relatively healthy versus subhealth with acute conditions or chronic diseases [81]); (2) the study methodology, comprising ROB and follow-up time frame; and (3) the intervention protocol, including PS dose, RT intensity (i.e., percentage of one repetition maximum), and treatment time period. The analyses results of NMR were expressed as β with 95% credible interval (CrI).

The participant compliance in response to PS plus RT was measured by all-cause withdrawal rate. In addition, any adverse event reported by the included RCTs was extracted and the corresponded occurrence rate was calculated to evaluate the adverse effects. The analyses results were presented as odds ratios (ORs) alongside 95% CIs.

All NMA and NMR analyses were performed using R statistical software (version 4.2.3, R Foundation for Statistical Computing, Vienna, Austria) [80,82]. For all the analyses, the statistical significance was accepted by a two-tailed *p* value of <0.05 .

2.8. Certainty of Evidence

The Grading of Recommendation Assessment, Development, and Evaluation (GRADE) system was employed to assess the quality of evidence contributing to NMA estimates for all main outcomes [83]. The GRADE framework evaluates five domains of each treatment arm derived from the included RCTs, namely study limitations, inconsistency, imprecision, incoherence, and publication bias. Based on the ranked score of each domain which was considered to upgrade or downgrade the quality of evidence, the evidence certainty was overall graded as high, moderate, low, and very low. The ranking and grading procedures were independently performed by two individuals (CDL, SWH) and validated by the others (HCC, MHH, THL, CLL).

We assessed the weighted contribution of direct comparisons to the NMA estimates for each outcome using the contributions matrix [84,85]. The downgrading level for the study limitation was based on the amounts of contribution of each direct estimate and corresponding risk of bias to the NMA estimates [86,87].


3. Results

3.1. Trial Selection Flowchart

The PRISMA flowchart of the trial selection process was presented in [Figure 1](#). Initially, a total of 1079 articles were identified by electronic searches and manual screen of literature. After excluding duplicates, we examining the retrieved titles of 471 studies and reviewed the abstracts of to assess their eligibility; followed by the full-text assessment of 126 relevant articles. A total of 78 registered clinical RCTs, corresponding with 85 published studies

[[88,89,90,91,92,93,94,95,96,97,98,99,100,101,102,103,104,105,106,107,108,109,110,111,112,113,114,115,116,117,118,119,120,121,122,123,124,125,126,127,128,129,130,131,132,133,134,135,136,137,138,139,140,141,142,143,144,145,146,147,148,149,150,151,152,153,154,155,156,157,158,159,160,161,162,163,164,165,166,167,168,169,170,171,172](#)], were included in this NMA and all of which were published between 1994 and 2023. The following included studies shared a common trial protocol respectively: 2 studies [[89,90](#)], 2 studies [[94,110](#)], 2 studies [[107,160](#)], 2 studies [[120,121](#)], 2 studies [[127,128](#)], 2 studies [[130,131](#)], and 2 studies [[147,148](#)].

Figure 1.



[Open in a new tab](#)

RRISMA flow diagram of study selection. PRISMA, Preferred Reporting Items for Systematic Reviews and Meta-Analyses.

3.2. Study Characteristics

[Table 1](#) summarizes the participant demographic data and study characteristics of the included RCTs. The details of each trial are presented in the [Supplementary Table S2](#). A total of 5272 participants were recruited among all included RCTs. In overall, the sample had a mean age of 70.2 (range: 47.0–89.2) years and a mean BMI of 25.8 (range: 18.6–33.3) kg/m². Among the 78 included RCTs ([Table S2](#)), 39 enrolled both women and men with an average proportion of female participants being 57.6% (range: 12.5–90.2%) whereas 22 had men-specific study design [[93,95,96,97,106,109,110,116,121,122,124,127,130,132,137,141,149,156,158,163,167,170](#)], 16 enrolled women only [[102,113,123,135,136,143,145,146,147,151,152,155,157,161,165,168](#)], and one did not report the sex distribution of the study sample [[103](#)].

Study Arm	Combined Treatment			Resistance Training (Combined with or without Placebo Supplement)			Protein Supplementation Alone		
	Trials (Groups), n ^a	Sample (n)	Mean (Range) b	Trials (Groups), n ^a	Sample (n)	Mean (Range) b	Trials (Groups), n ^a	Sample (n)	Mean (Range) b
Untrained healthy	33 (38)	947		27 (29)	692		8 (8)	202	
Dynapenia	4 (4)	92		3 (3)	78		2 (2)	39	
Frailty	12 (13)	394		12 (12)	356		2 (2)	44	
Sarcopenia	18 (20)	516		11 (11)	202		6 (6)	105	
Sedentary or inactive	7 (9)	213		5 (5)	100		2 (2)	25	
Overweight or obesity	5 (6)	296		5 (5)	236		0		
Protein source of supplementation									
Whey	43 (47)	1311					11 (11)	243	
Milk	17 (18)	553					2 (2)	73	
Soy	14 (14)	324					6 (6)	99	
Casein	4 (4)	94					0		
Meat	5 (5)	194					0		
Peanut	1 (1)	20					0		
Intervention compliance (%)									
Protein supplement	40 (47)	1287	91.5 (64–100)				37 (37)	323	87.8 (73–100)
Resistance training	46 (52)	1505	87.3 (44–100)	38 (38)	1117	86.2 (44–100)			
Placebo supplement				15 (15)	300	93.8 (71.5–100)			
Muscle mass (baseline)									
Total lean mass, kg	41 (47)	1354	45.9 (31.8–	36 (37)	1050	46.2 (32.0–	7 (7)	158	46.2 (39.9–

Study Arm	Combined Treatment		Resistance Training (Combined with or without Placebo Supplement)				Protein Supplementation Alone			
			Trials (Groups), n ^a	Sample (n)	Mean (Range) b	Trials (Groups), n ^a	Sample (n)	Mean (Range) b		
Item	Trials (Groups), n ^a	Sample (n)	Mean (Range) b	Trials (Groups), n ^a	Sample (n)	Mean (Range) b	Trials (Groups), n ^a	Sample (n)	Mean (Range) b	
			59.2)				60.8)			56.6)
Appendicular lean mass, kg	38 (41)	1160	18.8 (7.9–30.5)	36 (36)	955	18.8 (8.0–30.0)	9 (9)	161	18.8 (12.8–27.2)	
Total lean index, kg/m ²	44 (50)	1406	17.2 (13.6–22.9)	39 (40)	1143	17.4 (13.9–22.5)	8 (8)	168	17.2 (15.5–19.0)	
Appendicular lean index, kg/m ²	39 (41)	1150	7.0 (3.0–9.4)	35 (35)	935	7.1 (3.0–9.4)	11 (11)	208	7.1 (5.4–8.8)	
Handgrip strength (baseline), kg										
Sex-specific trial										
Men	7 (8)	282	30.3 (15.3–41.1)	4 (4)	64	33.2 (19.7–42.2)	2 (2)	37	25.5 (18.8–30.0)	
Women	4 (5)	93	20.2 (15.8–22.4)	4 (4)	71	20.3 (15.1–23.1)	1 (1)	25	23	
Sex mixed trial	23 (25)	819	26.5 (13.8–41.2)	19 (19)	619	26.8 (15.8–41.8)	6 (6)	119	22.3 (16.4–35.6)	
Physical mobility (baseline)										
5-time chair rise, s	28 (31)	1032	12.2 (4.6–26.5)	25 (25)	850	13.6 (4.7–30.0)	4 (4)	83	12.6 (7.7–18.7)	
Walking speed, m/s	36 (39)	1277	1.26 (0.44–2.33)	31 (31)	956	1.18 (0.51–2.05)	13 (13)	322	1.25 (0.45–2.14)	
Global mobility ^c	16 (16)	402	9.0 (6.2–	10 (10)	181	9.3 (7.1–	1 (1)	20	6.2	

Study Arm	Resistance Training (Combined with or without Placebo Supplement)			Protein Supplementation Alone		
	Trials (Groups), n ^a	Sample (n)	Mean (Range) b	Trials (Groups), n ^a	Sample (n)	Mean (Range) b
Item			11.4)			11.2)

[Open in a new tab](#)

^a Number of trials that reported the indicated item. ^b All summations calculated on the basis of the values reported in the analyzed studies that could be estimated. ^c The score is derived from the Short Physical Performance Battery.

With respective to participant physical conditions, 33 RCTs out of 78 [92,93,95,96,97,102,106,108,110,112,113,116,118,119,121,123,124,125,130,133,134,136,138,139,141,143,146,151,155,161,162,163,167,168] enrolled community-dwelling untrained middle-aged and older adults who were medically stable and relatively healthy, whereas others recruited individuals who were identified as sedentary or inactive (7 RCTs [91,103,109,126,129,149,153]), overweight or obesity (5 RCTs [140,156,159,164,170]), dynapenia (4 RCTs [88,98,104,169]), frailty (12 RCTs [101,105,111,114,117,132,142,147,150,152,157,160]), and sarcopenia (18 RCTs [89,99,100,115,122,127,135,137,144,145,154,158,165,166,169,171,172]). The study population were enrolled in America (29 RCTs), Asia (18 RCTs), Europe (23 RCTs), and Oceania areas (8 RCTs; Table 1 and Table S2).

Among all participants, 2458 (46.6%) received PS in combination with RT, 415 (7.9%) received PS alone, 1664 (31.6%) received RET alone, and 735 (13.9%) received RC (Table 1). With respect to the follow-up time points for outcome assessment, 57 out of 78 included RCTs conducted a short-term duration of ≤12 weeks; 23 out of 78 RCTs [91,98,99,100,101,102,110,112,123,126,134,136,137,140,143,144,150,151,152,158,160,164,172] employed a medium-term time frame ranging from 14 to 24 weeks, and 4 RCTs [127,130,139,144] assessed a long-term outcome ranging from 9 to 12 months (Table S2).

3.3. Protocol of Protein Supplementation

The characteristics for PS protocol are summarized in the Supplementary Table S2. In this NMA, a total of 6 types of protein supplement were identified (Table 1), namely whey protein (WP, 43 RCTs); milk protein (MP, 17 RCTs); soy protein (SP, 14 RCTs); casein (4 RCTs); beef meat (5 RCTs); and peanut (1 RCT). Regarding the PS protocol, 25 out of 78 included RCTs prescribed supplementation immediately after RT on the training days with a mean dosage of 26 (range 10–60) g/day; and the other 53 employed daily supplementation by a mean intake amount of 30 (range 10–80) g during an RT intervention (Table S2).


3.4. Protocol of Resistance Training

[Table S2](#) summaries the characteristics of RT protocols. In overall, the training load was moderate to high with a median (range) intensity of 75% (30–85%) of one-repetition maximum or 7 (5–9) of 10-point rating of perceived exertion. Regarding the training duration, 54 out of 78 included RCTs conducted a short intervention of 12 weeks or less, corresponding with 6–84 sessions; 21 RCTs employed a medium intervention of 14 weeks to 6 months, corresponding with 6–84 sessions; and 3 RCTs [[127](#),[130](#),[139](#)] had a long intervention up to 12 months.

3.5. Risks of Bias in Individual Study and Across Studies

[Figure 2](#) shows a summary of the risk-of-bias analysis for each domain and an overall rank crossing included RCTs. The details of each ranked risk-of-bias item per trial are presented in the [Supplementary Figure S1](#). Using the Cochrane risk-of-bias tool, over half of the included RCTs (43 out of 78, 55.1%) were ranked as having an overall high risk of bias whereas it determined 12 RCTs (15.4%) had low risk of bias and 23 RCTs (29.5%) had some concerns of bias ([Figure S1](#)). In total, the common domain of biases which obviously contributed to high risk of bias across all RCTs included at least one of the follows: selection, performance, detection, and attrition biases ([Figure 2](#)).

Figure 2.



[Open in a new tab](#)


Risk of bias graph crossing trials.

Risk of bias analysis crossing trials showed that 41 RCTs (52.6%) had a substantial unclear or high risk of selection bias raising from insufficient information regarding procedures of randomization or group assignment. Thirty-four RCTs out of 78 (43.6%) were considered having high risks of performance bias for blinding the participants and 27 RCTs (34.6%) were judged having high risks of detection bias because of the unblinded assessors. In addition, 40 RCTs (51.3%) which did not perform intention-to-treat analyses were scored as high risks of attrition bias; and low risk of reporting bias was ranked among all the included trials. Regarding other bias, we assumed that use of PS products supported by any funds or grants may arise potential agenda bias and therefore information of funding sources and conflict-of-interest declaration were assessed. There was potentially high risk of agenda bias in 11 RCTs (14.1%) which reported funding sources without any conflict-of-interest disclosures whereas 6 RCTs (7.7%) which did not report any information regarding funding sources and conflict of interest were considered unclear risk of agenda bias ([Figure 2](#)).

3.6. Treatment Efficacy for Muscle Mass

[Figure 3](#) presents the network connections among all treatment arms for muscle mass outcome. A total of 11 treatment options were identified in this NMA, included 6 combined treatments namely WP plus RT (WP + RT), MP plus RT (MP + RT), casein plus RT (Casein + RT), meat plus RT (Meat + RT), SP plus RT (SP + RT), peanut plus RT (Peanut + RT), 5 monotherapies namely WP, MP, SP, meat, and RT.

Figure 3.



[Open in a new tab](#)

Network diagram illustrating the interconnections among different treatment arms for muscle mass. Each node represents a treatment arm, and the size of the node is proportional to the number of participants assigned to the specific treatment. The thickness of each line is proportional to the number of direct comparisons between arms denoted on the line. RT, resistance training.

3.6.1. Conventional Pairwise Meta-Analysis


Muscle mass outcome were reported by 73 RCTs along with 12 treatment options and 151 pairwise comparisons in NMA (Figure 3). Direct comparisons of pairwise meta-analyses indicated that WP + RT (SMD = 1.36, 95% CI: 0.90–1.83), MP + RT (SMD = 1.31, 95% CI: 0.41–2.22), SP + RT (SMD = 1.06, 95% CI: 0.49–1.63) and RT (SMD = 0.79, 95% CI: 0.42–1.15) were more efficacious than RC for increasing muscle mass (Supplementary Table S3). In addition, the combined treatments WP + RT (SMD = 0.95, 95% CI: 0.52–1.37) and SP + RT (SMD

= 1.21, 95% CI: 0.58–1.85) achieved favor effects on muscle mass gain compared with its monotherapy of WP and SP, respectively. While comparing with RT alone, similar results were observed in WP + RT (SMD = 0.55, 95% CI: 0.17–0.93) and WP + RT (SMD = 0.56, 95% CI: 0.33–0.79).

3.6.2. Network Meta-Analysis

Considering RC as reference, the combined treatments namely WP + RT (SMD = 1.29), MP + RT (SMD = 1.24), Meat + RT (SMD = 1.23), and SP + RT (SMD = 1.06) achieved favorable outcomes for muscle mass increase regardless of follow-up time ([Figure 4](#)). Among all treatment arms, WP + RT was ranked as the most effective (SUCRA = 0.86) for muscle mass gain—followed by MP + RT (SUCRA = 0.83), and Meat + RT (SUCRA = 0.80) ([Figure 4](#)). The NMA results were based on significant global heterogeneity across treatment arms ($\tau^2 = 0.36$, $I^2 = 73.4\%$, $p < 0.0001$). Q statistic results assessing total consistency between designs in NMA showed insignificant under the assumption of a full design-by-treatment random-effects model ($\tau^2 = 0.35$, $p > 0.05$). According to the node-splitting analyses results, there was no inconsistencies between direct and indirect effects in NMA for muscle mass gain ([Figure S2](#)).

Figure 4.



[Open in a new tab](#)

Relative efficacy of treatment regimens on muscle mass gain during an overall follow-up duration. Each point estimate (square) presents the network combined effect (SMD) on the outcome measure relative to RC, with the 95% CI (horizontal line). SMD, standardized mean difference; CI, confidence interval; SUCRA, surface under the cumulative ranking curve; RT, resistance training; RC, regular care.

3.6.3. Subgroup Analysis Based on Follow-Up Time


The combined treatment WP + RT yielded the most effective treatment for muscle mass (SMD = 1.32, SUCRA = 0.83) over the short-term duration, whereas MP + RT was ranked the highest probabilities of successful treatment over the medium-term (SMD = 1.48, SUCRA = 0.93) and long-term (SMD = 1.14, SUCRA = 0.83) durations ([Figure S3](#)).

3.7. Treatment Efficacy for Muscle Strength

Muscle strength outcomes were assessed in terms of handgrip strength and leg strength. A total of 37 RCTs reported treatment outcome of handgrip strength along with 9 treatment regiments and 78 pairwise

comparisons in NMA ([Figure 5A](#)). The treatment effects on leg strength were investigated by 56 RCTs along with 11 treatment regiments and 104 pairwise comparisons in NMA ([Figure 5B](#)).

Figure 5.



[Open in a new tab](#)

Network diagram illustrating the interconnections among different treatment arms for muscle strength. Each node represents a treatment arm, and the size of the node is proportional to the number of participants assigned to the specific treatment. The thickness of each line is proportional to the number of direct comparisons between arms denoted on the line. RT, resistance training.

3.7.1. Conventional Pairwise Meta-Analysis


Results of direct comparisons indicated that PS plus RT achieved favorable effects on handgrip strength gain compared to RC ([Supplementary Table S4](#)), particularly the protein types of whey (SMD = 1.53, 95% CI: 0.88–2.19), casein (SMD = 2.98, 95% CI: 0.99–4.97), and soy (SMD = 1.51, 95% CI: 0.11–2.90). In addition, WP + RT yielded greater changes in handgrip strength than WP alone did (SMD = 0.86, 95% CI: 0.11–1.61); similar results were presented for the comparison between SP + RT and SP alone (SMD = 2.46, 95% CI: 0.44–4.48; [Table S4](#)).

The pairwise meta-analysis results for leg strength showed that WP + RT (SMD = 1.30, 95% CI: 0.73–1.87) and SP + RT (SMD = 1.73, 95% CI: 1.14–2.32) achieved favorable effects on leg strength gain compared to RC ([Supplementary Table S5](#)). Additionally, the combined treatments WP + RT yielded greater changes in leg strength than the monotherapy WP (SMD = 1.30, 95% CI: 0.76–1.84) or RT (SMD = 0.45, 95% CI: 0.20–0.69) did; similar results were observed in SP + RT when it compared with SP alone (SMD = 1.92, 95% CI: 1.16–2.68; [Table S5](#)).

3.7.2. Network Meta-Analysis

There were significant effects in favor of PS plus RT for handgrip strength compared to RC, particularly the protein sources derived from whey (SMD = 1.46), soy (SMD = 1.41), and milk (SMD = 1.27), during an overall follow-up duration (Figure 6A). Among all treatment options, WP + RT yielded the highest probabilities of treatment effects (SUCRA = 0.84)—followed by SP + RT (SUCRA = 0.78) and MP + RT (SUCRA = 0.72)—during the overall follow-up duration (Figure 6A). There was significant global heterogeneity across all treatment arms involved in NMA model ($\tau^2 = 0.88$, $I^2 = 91.7\%$, $p < 0.0001$). Q statistic results assessing total consistency between designs in NMA showed insignificant under the assumption of a full design-by-treatment random-effects model ($\tau^2 = 0.88$, $p > 0.05$). According to the node-splitting analyses results, there was no inconsistencies between direct and indirect effects in NMA for handgrip strength gain (Figure S4).

Figure 6.



[Open in a new tab](#)

Relative efficacy of treatment regimens on muscle strength during an overall follow-up duration. Each point estimate (square) presents the network combined effect (SMD) on the outcome measure relative to RC, with the 95% CI (horizontal line). SMD, standardized mean difference; CI, confidence interval; SUCRA, surface under the cumulative ranking curve; RT, resistance training; RC, regular care.

With respect to treatment outcome of muscle strength in the lower extremity (Figure 6B), majority of the PS regimens augmented increases in leg strength during RT, including whey, milk, soy, and meat (SMD = 1.27–1.41). In addition, WP + RT yielded the most effective (SUCRA = 0.87) for muscle strength in the legs—followed by MP + RT (SUCRA = 0.80) and SP + RT (SUCRA = 0.79; Figure 6B). Significant global heterogeneity was observed ($\tau^2 = 0.30$, $I^2 = 75.6\%$, $p < 0.0001$). Q statistic results assessing total consistency between designs in NMA showed insignificant under the assumption of a full design-by-treatment random-effects model ($\tau^2 = 0.88$, $p > 0.05$). According to the node-splitting analyses results, there was no inconsistencies between direct and indirect effects in NMA for leg strength (Figure S5).

3.7.3. Subgroup Analysis Based on Follow-Up Time


In handgrip strength, WP + RT yielded the highest medium-term (SMD = 1.74, SUCRA = 0.83) and long-term (SMD = 1.69, SUCRA = 0.99) treatment efficacy, respectively ([Figure S6](#)). Within the short-term time frame, SP + RT was ranked the most effective among whole treatment regimens in NMA (SMD = 1.42, SUCRA = 0.91).

For leg strength gain, the combined regimen SP + RT yielded the highest short-term efficacy (SMD = 1.52, SUCRA = 0.93), whereas MP + RT was ranked the most effective among all treatment regimens over the medium-term (SMD = 1.11, SUCRA = 0.95) and long-term (SMD = 1.21, SUCRA = 0.95) time frames, respectively ([Figure S7](#)).

3.8. Effectiveness of Treatment for Physical Mobility

Treatment efficacy on physical mobility were assessed by walking speed (35 RCTs), chair rise (31 RCTs), TUG (8 RCTs), and global function in terms of SPPB score (13 RCTs). In walking speed, a total of 9 treatment options were identified along with 91 pairwise comparisons in NMA ([Figure 7A](#)). In addition, a total of 8 treatment options (62 pairwise comparisons) and 6 treatment options (12 pairwise comparisons) were identified in NMA for chair rise ([Figure 7B](#)) and TUG ([Figure 7C](#)), respectively. For SPPB, 6 treatment options were identified along with 18 pairwise comparisons in NMA ([Figure 7D](#)).

Figure 7.



[Open in a new tab](#)

Network diagram illustrating the interconnections among different treatment arms for physical mobility. Each node represents a treatment arm, and the size of the node is proportional to the number of participants assigned to the specific treatment. The thickness of each line is proportional to the number of direct comparisons between arms denoted on the line. RT, resistance training; SPPB, Short Physical Performance Battery.

3.8.1. Conventional Pairwise Meta-Analysis


Direct pairwise comparisons indicated that WP + RT achieved favorable effects on walking-speed recovery (SMD = 0.84, 95% CI: 0.43–1.25; [Supplementary Table S6](#)), chair rise (SMD = 0.85, 95% CI: 0.46–1.25; [Supplementary Table S7](#)), and SPPB score increase (SMD = 1.79, 95% CI: 1.09–2.48; [Supplementary Table S8](#)) compared to RC, respectively. The monotherapy WP (SMD = 0.76, 95% CI: 0.30–1.22; [Table S6](#)) achieved

favorable effects on walking-speed recovery compared to RC, as well as RT did in walking speed (SMD = 0.59, 95% CI: 0.22–0.95; [Table S6](#)) and chair rise (SMD = 0.82, 95% CI: 0.53–1.11; [Table S7](#)). In TUG, direct pairwise comparison results showed no significant effects among all treatment options with reference to RC ([Table S9](#)).

3.8.2. Network Meta-Analysis

For walking speed recovery during an overall time ([Figure 8A](#) and [Figure S8](#)), the composed regimens WP + RT (SMD = 0.73) and MP + RT (SMD = 0.53) achieved favorable effects compared to RC. Among all treatment options, WP + RT was ranked the most effective treatment to increase walking speed (SUCRA = 0.84; [Figure S8](#)). Significant global heterogeneity was observed ($\tau^2 = 0.28$, $I^2 = 58.9\%$, $p < 0.0001$). The full design-by-treatment random-effects model determined that the overall inconsistency between designs in NMA was significant ($\tau^2 = 0.29$, $p < 0.0001$). According to the node-splitting analyses results, it showed no inconsistencies between the direct and indirect comparisons in NMA ([Figure S9](#)).

Figure 8.



[Open in a new tab](#)

Relative efficacy among treatment regimens on changes in physical mobility scores during an overall study time. Each point estimate (square) presents the network combined effect on the specific outcome measure, with the 95% CI (horizontal line). The positive results indicate favorable efficacy of the treatment regimen compared to RC. The blue-colored point represents the highest probability of effects among all treatment regimens. Std, standardized; CI, confidence interval; RT, resistance training; SPPB, Short Physical Performance Battery; RC, regular care.

For chair-rise performance at an overall time frame ([Figure 8B](#) and [Figure S10](#)), the NMA results showed favorable effects responding to WP + RT (SMD = 0.84), as well as SP + RT (SMD = 0.44) and MP + RT (SMD = 0.31), compared to RC. Of all treatment regimens, WP + RT was ranked the optimal treatment to increase

chair-rise ability (SUCRA = 0.88; [Figure S10](#)). Significant global heterogeneity of the NMA model was identified ($\tau^2 = 0.12$, $I^2 = 57.0\%$, $p < 0.0001$). Using a full design-by-treatment random-effects model, it determined an insignificant overall between-design inconsistency of the NMA ($\tau^2 = 0.16$, $p > 0.05$). The node-splitting analyses did not indicate any inconsistency between direct and indirect effects in NMA ([Figure S11](#)).

Regarding TUG task, the NMA results showed no favorable effects responding to any of treatment arms ([Figure 8C](#) and [Figure S12](#)). Among all treatment arms, MP + RT was ranked as the most effective treatment to save time in completing the TUG task (SUCRA = 0.79; [Figure S12](#)). No evidence of the global heterogeneity was identified ($\tau^2 < 0.0001$, $I^2 = 0\%$, $p > 0.05$). The overall between-design inconsistency of the NMA showed insignificant ($\tau^2 = 0.005$, $p > 0.05$). Based on the node-splitting analyses results, there was no inconsistencies between direct and indirect evidence ([Figure S13](#)).

With respect to treatment outcome for global function, in terms of SPPB score, the NMA results showed that WP (SMD = 1.45), as well as MP (SMD = 0.97), in combination with RT obtained favorable effects on SPPB score relative to RC, regardless of study time frame ([Figure 8D](#) and [Supplementary Figure S14](#)). Of all treatment regimens, WP + RT was ranked the optimal treatment to increase SPPB score (SUCRA = 0.87; [Figure S14](#)). There was significant global heterogeneity of the NMA model for SPPB ($\tau^2 = 0.27$, $I^2 = 75.2\%$, $p < 0.0001$). There was significant overall between-design inconsistency ($\tau^2 = 0$, $p < 0.0001$). The node-splitting analyses did not indicate any inconsistency between direct and indirect comparisons ([Figure S15](#)).

3.8.3. Subgroup Analysis Based on Follow-Up Time

During the short-term time, WP + RT yielded the optimal treatment option to increase performances in walking speed (SMD = 0.74, SUCRA = 0.83; [Figure S8](#)) and chair rise (SMD = 0.98, SUCRA = 0.85; [Figure S10](#)), as well as MP + RT did for saving timed-up-and-go time (SMD = 0.33, SUCRA = 0.76; [Figure S12](#)) and increasing total score of SPPB (SMD = 0.45, SUCRA = 0.71; [Figure S14](#)).

Within the medium-term follow-up time frame, WP + RT was ranked the most effective treatment option for walking-speed recovery (SMD = 0.25, SUCRA = 0.78; [Figure S8](#)) and to increase total score of SPPB (SMD = 2.19, SUCRA = 0.88; [Figure S14](#)), as well as SP + RT and MP + RT did for increasing chair-rise ability (SMD = 0.87, SUCRA = 0.88; [Figure S10](#)) and saving TUG time (SMD = 0.33, SUCRA = 0.76; [Figure S12](#)), respectively.

Regarding the long-term treatment outcome, WP + RT was ranked the most effective treatment option for walking-speed recovery (SMD = 0.58, SUCRA = 0.98; [Figure S8](#)). None of the included RCTs conducted long-term follow up for chair rise, TUG, and SPPB outcome.

3.9. Network Meta-Regression Analyses


Effects of substantial moderators for each main outcome are presented in the [Supplementary Table S10](#). Health status had effects on relative treatment efficacy among treatment arms. Participants who suffered acute (hospitalized) or chronic (obese, sarcopenic, frail, and mobility limited) conditions may achieved greater effects on muscle-mass changes ($\beta = 0.74$; 95% CrI: 0.22, 1.25) compared with those who were relatively healthy. In

addition, there was potential gender effect on relative treatment efficacy for muscle strength and physical function. A greater proportion of women in the study sample predicted minor effects on handgrip strength ($\beta = -1.72$; 95% CrI: $-2.68, -0.77$) and SPPB ($\beta = -1.61$; 95% CrI: $-2.47, -0.79$). Other potential moderators of relative treatment efficacy were identified, including amounts of PS ($\beta = 0.76$; 95% CrI: $0.06, 1.47$) and BMI ($\beta = -1.85$; 95% CrI: $-2.89, -0.81$) for leg strength and SPPB, respectively. In addition, a longer intervention period ($\beta = 1.41$; 95% CrI: $0.78, 2.05$), as well as follow-up time ($\beta = 1.41$; 95% CrI: $0.77, 2.03$), was associated with greater increases in SPPB score. Furthermore, area of study population had significant influences in strength gains of handgrip ($\beta = -1.04$; 95% CrI: $-2.00, -0.07$) and leg extensors ($\beta = -0.86$; 95% CrI: $-1.70, -0.05$). No moderator was identified to have an influence on the treatment efficacy regarding mobility tasks of walking speed, chair rise, and TUG.

3.10. Compliance and Side Effects

As a whole, an all-cause attrition rate of 0–78.6% was reported among the included RCTs, of which 0% to 45.2% were dropout due to treatment noncompliance ([Table S11](#)). Eleven RCTs reported withdraw of participants due to noncompliance of PS by an attrition rate of 2.0–32.1% [[89,91,103,113,119,126,136,143,153,159,169](#)]. Noncompliance of RT were reported by 17 RCTs with an attrition rate of 4.5–31.3% [[91,101,103,111,114,121,124,125,126,133,138,140,151,159,164,169,170](#)]. There was no difference in compliance across all the treatment regimens with reference to RC ([Figure 9A](#)).

Figure 9.



[Open in a new tab](#)

Participant compliance and side effects of treatment regimens. Data concerning treatment-related (A) withdrawals and (B) adverse events were pooled using inverse variance weighting methods. Each point estimate (square) presents the network combined effect (OR) of the indicated treatment arm relative to RC, with the 95% CI (horizontal line). OR, odds ratio; CI, confidence interval; RT, resistance training.

There was no serious adverse event occurred after PS plus RT or its monotherapy. In total, 12 [89,97,98,108,111,117,125,126,136,140,153,159] and 15 [97,98,102,111,112,116,117,121,124,126,130,140,159,164,166] out of the 78 included RCTs reported mild adverse events in response to PS and RT, respectively. Majority of the reported adverse events were RT-induced joint pain and muscle soreness (Table S11) and gastrointestinal symptoms such as diarrhea, bloating, and constipation after PS. Compared with RC, there was no significant adverse effects among all treatment regimens excepting WP + RT (Figure 9B).

3.11. Publication Bias

Publication biases among the included RCTs for all outcome domains are presented in the Supplementary Figure S16. The funnel plot indicated no substantial asymmetry of publication bias across the included RCTs. Begg–Mazumdar test results for muscle mass ($p < 0.01$; Figure S16A) and leg muscle strength ($p < 0.01$; Figure S16C) indicated obvious reporting bias among the RCTs included in the NMA. No significant bias was observed in other main outcomes.

3.12. Certainty of the Evidence

[Table 2](#) summarizes the GRADE certainty rating for all main outcomes and the details of judgments in each domain of GRADE framework are presented in Supplementary [Tables S12–S18](#). In overall, certainty of the evidence ranged from low to high among combined treatments for muscle mass, strength, and mobility outcomes, particularly WP + RT, SP + RT, MP + RT, and Meat + RT ([Table 2](#)). The most common judgements downgrading the certainty of evidence were associated with major concerns about study limitation, imprecision, and publication bias.

Table 2.

GRADE certainty rating of treatment options in all main outcomes.

Treatment (Common Comparator: Regular Care)	GRADE Certainty of Evidence ^a						
	Muscle Mass	Muscle Strength		Physical Mobility			
		Handgrip	Leg Strength	Walking Speed	Chair Rise	Timed Up and Go	SPPB
Combined therapy							
Casein + RT	⊕⊖⊖⊖ b,d,e	⊕⊖⊖⊖ b,c,d	⊕⊖⊖⊖ b,d,e				⊕⊕⊖⊖ b,e
Meat + RT	⊕⊕⊖⊖ b,e	⊕⊕⊖⊖ b,d	⊕⊕⊖⊖ b,e	⊕⊕⊖⊖ b,d	⊕⊕⊖⊖ b,d	⊕⊕⊖⊖ b,d	
Milk + RT	⊕⊕⊖⊖ b,e	⊕⊕⊕⊖ b	⊕⊕⊖⊖ b,e	⊕⊕⊕⊖ b	⊕⊕⊕⊖ b	⊕⊕⊖⊖ b,d	⊕⊕⊕⊖ b
Peanut + RT	⊕⊖⊖⊖ b,d,e						
Soy + RT	⊕⊕⊖⊖ b,e	⊕⊕⊕⊕	⊕⊖⊖⊖ b,c,e	⊕⊕⊖⊖ b,d	⊕⊕⊕⊕	⊕⊕⊖⊖ b,d	
Whey + RT	⊕⊕⊖⊖ b,e	⊕⊕⊕⊖ b	⊕⊕⊖⊖ b,e	⊕⊕⊕⊖ b	⊕⊕⊕⊖ b	⊕⊕⊖⊖ b,d	⊕⊕⊖⊖ b,c
Monotherapy							
Meat	⊕⊖⊖⊖ b,d,e		⊕⊖⊖⊖ b,d,e				
Milk	⊕⊖⊖⊖ b,d,e		⊕⊖⊖⊖ b,d,e	⊕⊕⊖⊖ b,d	⊕⊕⊖⊖ b,d		
Soy	⊕⊖⊖⊖ b,d,e	⊕⊕⊕⊖ d	⊕⊖⊖⊖ b,c,d,e	⊕⊕⊖⊖ b,d			
Whey	⊕⊖⊖⊖ b,d,e	⊕⊕⊖⊖ b,d	⊕⊖⊖⊖ b,d,e	⊕⊕⊖⊖ b,c	⊕⊕⊖⊖ b,d		⊕⊕⊕⊖ b
Resistance training	⊕⊕⊖⊖ b,e	⊕⊕⊕⊖ b	⊕⊕⊖⊖ b,e	⊕⊕⊕⊖ b	⊕⊕⊕⊖ b	⊕⊕⊖⊖ b,d	⊕⊕⊕⊖ b

^a Certainty of evidence is graded as follows: High: ⊕⊕⊕⊕; Moderate: ⊕⊕⊕⊖; Low: ⊕⊕⊖⊖; Very low: ⊕⊖⊖⊖. ^b Network contribution of high risk of bias ≥ 50%. ^c There is significant difference between direct and indirect estimates. ^d 95% confidence interval is wide and imprecise. ^e Test for publication bias is statistically significant. GRADE, Grading of Recommendations, Assessment, Development and Evaluations; SPPB, Short Physical Performance Battery.

4. Discussion

The main purpose of this study was to identify the comparative efficacy of different protein sources combined with RT for muscle mass, strength, and physical mobility outcomes in middle-aged and older adults. The main results showed that (1) the combined treatment of protein supplement and RT (particularly WP + RT, SP + RT, MP + RT, and Meat + RT) exerted favorable effects on muscle mass and strength gain relative to RC, whereas WP + RT and MP + RT achieved promising effects on physical mobilities including walking speed, chair rise, and SPPB; (2) based on the cumulative ranking results, whey protein supplements combined with RT yielded the most effective treatment to increase muscle mass, strength, and physical mobility, respectively; (3) a number of potential moderators of relative treatment efficacy were identified, including BMI, gender, healthy status, study area, supplement dosage, and treatment time. In addition, the composite PS plus RT exhibited high compliance despite of higher occurrence rate of nonserious adverse events among PS regimens, particularly WP + RT.

A number of systematic reviews and NMAs have compared the effects among different protein sources of supplementation during exercise training in young and older people [44,62,63,64,65,66,67], and results have generally indicated no significant difference in lean mass or strength gains between WP versus SP [62,63,66], WP versus MP [44], WP versus beef [64], and SP versus other mixed protein sources [63] during an exercise intervention. However, more other protein sources, such as casein and peanut, have yet to be comprehensively compared with conventional protein supplements. In the present study which targeted PS during an RT intervention in middle-aged and older populations, six protein sources (i.e., whey, soy, milk, meat, casein, and peanut) were identified and fully compared in an NMA. Our analyzed results are generally in agreement with those of the previous systematic reviews. We further rank the probabilities of successful treatment among multiple regimens of PS, RT, and its combinations, and the results showed that WP + RT yielded the most effective treatment to increase muscle mass and strength, and to enhance mobility performance. Additionally, the present results indicate that combined treatments incorporating PS and RT can yield extra benefits compared with its monotherapies in adults who aged ≥ 45 years.

In the present study, the ranked results indicated that WP yielded the most effective supplement to augment RT-induced muscle mass gain, followed by MP, meat, SP, and casein (Figure 4). Our findings were supported by previous studies. First, PS monotherapy by WP supplements achieves greater changes in postprandial muscle protein synthesis rate compared with other protein sources, especially in older individuals [67,173], and such favorable effects become more significant for WP + RT [67]. Reasonably, it is expected that the superior effects of WP supplementation on postprandial muscle protein synthesis may lead to favorable long-term changes in RT-induced skeletal muscle adaptation. Secondly, the postprandial muscle protein synthetic response to the ingestion of different protein sources have been compared under RT intervention [61,174,175,176]. The

previous results revealed that WP supplements stimulate higher postprandial muscle protein synthesis rates than SP [175] and Casein [61,175] do. In addition, the ingestion of SP after RT stimulates muscle protein synthesis to a greater extent when compared with casein [175], as well as the MP ingestion does while comparing with beef protein [174] and SP supplements [176], respectively. Accounting these previous results together, it suggests that WP exhibits similar effects with MP and is superior to MP, meat, SP, and casein with regard to stimulate muscle protein synthesis, which further supports the ranked results among multiple protein sources identified in this NMA regarding treatment efficacy on muscle mass gain.

Based on the NMA results in the current study, the protein source WP was ranked as the optimal regimen for muscle mass gain during RT intervention and such superiority in treatment efficacy was simultaneously observed in strength and physical mobility. Our findings were in agreement with previous meta-analyses which have concluded that PS synergistically increases muscle mass along with strength gain and physical mobility restoration in older people undergoing exercise training enhancement [177,178,179,180]. Given the facts that muscle mass changes in response to PS significantly contribute to additional increases in strength and walking capability during exercise training [181] and that increased leg strength is associated with faster habitual walking speed following PS plus RT [88], especially in those at high risk of sarcopenia and frailty, the superior effects of WP + RT on muscle mass may parallelly give rise to the superiority of effects on strength and mobility while comparing with other protein sources in this NMA.

In this NMA, a number of substantial moderators were identified to have significant effects on relative efficacy among treatment regimens. First, greater treatment effects on muscle mass were likely to be observed in the participants with acute conditions (hospitalized) or chronic diseases (obese, sarcopenic, frail, and mobility limited) compared with their relatively healthy peers (Table S10). Such findings are in line with the previous systematic review studies indicating that individuals with sarcopenia or frailty risks tend to obtain greater lean mass gains after PS plus exercise training than their healthy peers do [182,183]. Comparing with healthy older, the individuals who suffered frailty, sarcopenia, or other comorbidities have experienced minor habitual intake of protein and lower physical activity level [184,185,186,187], which impacts myofibrillar protein synthetic rates following PS and RT [52,53,187]. However, the ingestion of additional protein during RT may contribute relatively more responses of myofibrillar protein synthesis in those with aging-related conditions than in normal control [188]. Under such scenario, older people with acute or chronic conditions are likely to benefit more in lean mass gains from PS plus RT compared with their healthy peers who achieve minor changes in muscle mass following PS plus RT [189]. Secondly, increases in proportion of women of the study sample may result in decreased changes in handgrip strength and SPPB following PS plus RT; such findings indicated substantial gender effects on muscular adaptations in response to PS plus RT, particularly for muscle strength and physical mobility. Our results are supported by the previous studies demonstrating sex-specific difference in treatment efficacy of PS plus exercise for muscle strength [119,152,190] and mobility (CR) [119] among older adults. Thirdly, the current NMR results showed positive associations between amount of protein supplement and leg strength gain, which indicated that PS induced augmentations of training effects on muscle strength might be dose dependent. Numeral previous studies have been observed that post-exercise increases in myofibrillar fractional synthesis rates responding to PS stimuli are significantly enhanced by an increased amount of protein intake in trained young or older adults [191,192,193]. Accordingly, such acute dose-dependent effects on muscle synthesis may lead to long-term neuromuscular adaptations following RT, particularly in terms of strength gain. In addition, our findings were agreed by the previous study concluding that ingestion of dietary protein as high

as 1.6 g/kg/day significantly enhances leg extension strength compared with the minor intake (0.8 g/kg/day) in elderly men [194]. Finally, we observed that a higher BMI was corresponded with minor changes in SPPB score whereas greater changes in SPPB score may be observed over a longer intervention or follow-up length up to 6 months. According to the present results, greater function recovery may occur at 6-month follow up, compared with that within a follow-up time of 3 months or shorter in older individuals who are not overweight or obese. Therefore, the restoration of physical mobility in response to PS plus RT appears to be time-dependent, especially for those who have normal body weight.

Participant compliance with PS was equally comparable among protocols of different protein sources in this NMA. The results indicated that older individuals tolerated well for all protein products derived from different sources. However, a relatively higher risk of occurrence rate of adverse events (OR = 3.04) in favor of WP was noted, comparing with that of other protein sources (OR = 1.13–2.04; Figure 9B). After consuming whey supplements, 45 adverse events in 45 participants were reported by 6 RCTs [89,97,98,108,126,140], most of which were related to gastrointestinal problems such as gastro esophageal reflux, nausea, bloating or diarrhea. Among the 45 participants who suffered WP-induced complications, 30 (66.7%) experienced mild to moderate severity of gastrointestinal distress and continued the supplementation by modifying dose, whereas the others (33.3%) happened severe gastrointestinal problems which needed to cease consuming whey supplements. Based on the results, we assume that most of the adverse events responding to WP are considered as nonserious and can be well treated despite of its higher occurrence rate.

Several limitations of the present NMA are needed to be noted. First, this NMA is limited by the varied study methodological quality with only 12 RCTs (15.4%) were ranked as low risk of bias, which restricts the strength of the results, and may lead to overestimation of relative effects among treatment regimens. Secondly, due to wide variation in PS (protein source, dosage, intake time) and RT protocols (training mode, duration, and volume) among included RCTs, it was difficult to make a definite protocol of specific regimen for treatment efficacy, even though WP + RT was ranked as the highest probability of treatment efficacy among all treatment arms. In addition, the optimal dosage of supplementation and training duration were not determined in the current NMA in spite of the fact that an increase of PS dose is associated with greater leg strength gain and a longer intervention is accompanied with greater effects on SPPB. Further researches are warranted to identify the optimal intake amount of protein supplements and training period, especially the WP + RT, for elder populations. Thirdly, significant publication biases of treatment effects were detected in muscle mass and leg strength, which downgrades the certainty of evidence. Therefore, the conclusions in these outcomes should be cautiously interpreted. Finally, some of the included RCTs conducted placebo supplementation in combination with RT or not. The placebo effects were not excluded for accounting the treatment effects of RT alone or RC groups. Therefore, the comparative effects among all treatment arms may be underestimated, as using the RC groups as the common reference.

5. Conclusions

The present NMA determined the comparative efficacy among different sources of PS for community-dwelling, hospitalized or institutionalized older individuals undergoing RT. Overall, the combined treatments of PS and RT were superior to its monotherapy. In addition, WP + RT was determined to be the optimal treatment regimen for

muscle mass and strength gains as well as restoration of physical mobility with large effects and low to moderate certainty of evidence, despite of its relatively high risk of nonserious side effects. The treatment efficacy appears to be mediated by healthy status, sex, and body weight, and is likely to be dose and time dependent particularly for restorations of strength and physical mobility. Our findings add the knowledge on RS and RT intervention strategies, especially for older individuals with high sarcopenia and frailty risks. The results of this NMA may help the clinicians to prescribe the optimal PS protocol during RT intervention for ensuring optimal treatment outcomes.

Acknowledgments

This manuscript was edited by the professional editing.

Supplementary Materials

The following supporting information can be downloaded at:

<https://www.mdpi.com/article/10.3390/nu16070941/s1> , Table S1. Database search formulas; Table S2.

Summary of included study characteristics; Table S3. League table for pairwise and network meta-analysis of mean change in muscle mass from baseline; Table S4. League table for pairwise and network meta-analysis of mean change in handgrip strength from baseline; Table S5. League table for pairwise and network meta-analysis of mean change in leg strength from baseline; Table S6. League table for pairwise and network meta-analysis of mean change in walking speed from baseline; Table S7. League table for pairwise and network meta-analysis of mean change in chair rise from baseline; Table S8. League table for pairwise and network meta-analysis of mean change in global physical mobility (SPPB) from baseline; Table S9. League table for pairwise and network meta-analysis of mean change in timed up-and-go task from baseline; Table S10. Associations of moderators with treatment efficiency for main outcomes; Table S11. Summary of compliance & adverse events; Table S12. Judgments in each domain of GRADE ranking system for muscle mass outcome; Table S13. Judgments in each domain of GRADE framework for handgrip strength outcome; Table S14. Judgments in each domain of GRADE framework for leg strength outcome; Table S15. Judgments in each domain of GRADE framework for walking speed outcome; Table S16. Judgments in each domain of GRADE framework for chair rise outcome; Table S17. Judgments in each domain of GRADE framework for timed up-and-go outcome; Table S18. Judgments in each domain of GRADE framework for timed global mobility (SPPB) outcome; Figure S1. Risk of bias summary for each trial; Figure S2. Forest plot of node-splitting results for muscle mass; Figure S3. Relative effects among treatment regimens for muscle mass gain at each follow-up time frame; Figure S4. Forest plot of node-splitting results for handgrip strength; Figure S5. Forest plot of node-splitting results for leg strength; Figure S6. Relative effects among treatment regimens for handgrip strength at each follow-up time frame; Figure S7. Relative effects among treatment regimens for leg strength at each follow-up time frame; Figure S8. Relative effects among treatment regimens for walking speed at each follow-up time frame; Figure S9. Forest plot of node-splitting results for walking speed; Figure S10. Relative effects among treatment regimens for chair rise at each follow-up time frame; Figure S11. Forest plot of node-splitting results for chair rise; Figure S12. Relative effects among treatment regimens for timed up-and-go task at each follow-up time frame; Figure S13. Forest plot of node-splitting results for TUG; Figure S14. Relative effects among treatment regimens for Short Physical Performance Battery at each follow-up time frame; Figure S15. Forest plot of node-splitting results for SPPB; Figure S16.

Funnel plots of the intervention effects for (A) muscle mass, (B) handgrip strength, (C) leg muscle strength, (D) walk speed, (E) chair rise ability, (F) timed up and go, and (G) SPPB.

[nutrients-16-00941-s001.zip](#) ^(6MB, zip)

Author Contributions

The authors' responsibilities are outlined as follows—Conceptualization, C.-D.L., S.-W.H. and C.-L.L.; searched for and selected relevant studies, C.-D.L., S.-W.H. and C.-L.L.; data extraction, C.-D.L., S.-W.H. and C.-L.L.; validation, T.-H.L. and C.-L.L.; formal analysis, C.-D.L. and H.-C.C.; data curation, S.-W.H. and C.-L.L.; writing—original draft preparation, C.-D.L. and S.-W.H.; writing—review and editing, H.-C.C., M.-H.H. and S.-W.H.; visualization and supervision, T.-H.L. and S.-W.H. All authors have read and agreed to the published version of the manuscript.

Institutional Review Board Statement

Ethical review and approval were not applicable for this study, due to the research design of systemic review and network meta-analysis. This systemic review and network meta-analysis followed the guidelines the guidelines of the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) and the study protocol was registered at the PROSPERO database (registration number: CRD42023472942). All of the included randomized controlled trials obtained individual approvals from the respective local research ethics committee.

Informed Consent Statement

Not applicable. All of the included randomized controlled trials obtained individual informed consents from the respectively enrolled participants.

Data Availability Statement

Refer to [Supplementary Materials](#). Raw data available on request.

Conflicts of Interest

The authors declare no conflicts of interest.

Funding Statement

This study was funded by grants from Taipei Medical University-Shuang Ho Hospital, Ministry of Health and Welfare, Taiwan (grant no. 112TMU-SHH-35) and Taipei Medical University (grant no. TMU110-AE1-B10). The funding sources played no role in the design, implementation, data analysis, interpretation, or reporting of the study. The contents of this publication are solely the responsibility of the authors and do not necessarily represent the official view of the funding sources.

Footnotes

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

References

1. Chen L.K., Woo J., Assantachai P., Auyeung T.W., Chou M.Y., Iijima K., Jang H.C., Kang L., Kim M., Kim S., et al. Asian Working Group for Sarcopenia: 2019 Consensus Update on Sarcopenia Diagnosis and Treatment. *J. Am. Med. Dir. Assoc.* 2020;21:300–307.e302. doi: 10.1016/j.jamda.2019.12.012. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
2. Cruz-Jentoft A.J., Bahat G., Bauer J., Boirie Y., Bruyere O., Cederholm T., Cooper C., Landi F., Rolland Y., Sayer A.A., et al. Sarcopenia: Revised European consensus on definition and diagnosis. *Age Ageing.* 2019;48:16–31. doi: 10.1093/ageing/afy169. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
3. Jung H., Tanaka S., Kataoka S., Tanaka R. Association of sarcopenia, pre-sarcopenia, and dynapenia with the onset and progression of locomotive syndrome in Japanese older adults: A cross-sectional study. *J. Physiol. Anthropol.* 2023;42:16. doi: 10.1186/s40101-023-00334-3. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
4. Longobucco Y., Krumpoch S., Lauretani F., Angileri V., Sieber C., Marzetti E., Calvani R., Cherubini A., Landi F., Bernabei R., et al. Gait characteristics in community-dwelling older persons with low skeletal muscle mass and low physical performance. *Aging Clin. Exp. Res.* 2022;34:1563–1571. doi: 10.1007/s40520-021-02061-0. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
5. Rolland Y., Lauwers-Cances V., Cristini C., Abellan van Kan G., Janssen I., Morley J.E., Vellas B. Difficulties with physical function associated with obesity, sarcopenia, and sarcopenic-obesity in community-dwelling elderly women: The EPIDOS (EPIDemiologie de l'OSteoporose) Study. *Am. J. Clin. Nutr.* 2009;89:1895–1900. doi: 10.3945/ajcn.2008.26950. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
6. Cawthon P.M., Orwoll E.S., Peters K.E., Ensrud K.E., Cauley J.A., Kado D.M., Stefanick M.L., Shikany J.M., Strotmeyer E.S., Glynn N.W., et al. Strong Relation between Muscle Mass Determined by D3-creatine Dilution, Physical Performance and Incidence of Falls and Mobility Limitations in a Prospective

Cohort of Older Men. *J. Gerontol. Ser. A Biol. Sci. Med. Sci.* 2019;74:844–852. doi: 10.1093/gerona/gly129. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

7. Damluji A.A., Alfaraidhy M., AlHajri N., Rohant N.N., Kumar M., Al Malouf C., Bahrainy S., Ji Kwak M., Batchelor W.B., Forman D.E., et al. Sarcopenia and Cardiovascular Diseases. *Circulation*. 2023;147:1534–1553. doi: 10.1161/CIRCULATIONAHA.123.064071. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

8. Pickering M.E., Chapurlat R. Where Two Common Conditions of Aging Meet: Osteoarthritis and Sarcopenia. *Calcif. Tissue Int.* 2020;107:203–211. doi: 10.1007/s00223-020-00703-5. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

9. Shorter E., Sannicandro A.J., Poulet B., Goljanek-Whysall K. Skeletal Muscle Wasting and Its Relationship With Osteoarthritis: A Mini-Review of Mechanisms and Current Interventions. *Curr. Rheumatol. Rep.* 2019;21:40. doi: 10.1007/s11926-019-0839-4. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

10. Vlietstra L., Stebbings S., Meredith-Jones K., Abbott J.H., Treharne G.J., Waters D.L. Sarcopenia in osteoarthritis and rheumatoid arthritis: The association with self-reported fatigue, physical function and obesity. *PLoS ONE*. 2019;14:e0217462. doi: 10.1371/journal.pone.0217462. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

11. Torii M., Itaya T., Minamino H., Katsushima M., Fujita Y., Tanaka H., Oshima Y., Watanabe R., Ito H., Arai H., et al. Management of sarcopenia in patients with rheumatoid arthritis. *Mod. Rheumatol./Jpn. Rheum. Assoc.* 2023;33:435–440. doi: 10.1093/mr/roac095. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

12. Luo C., Liu R.Y., Zhang G.W., Hu F., Jin Y.H., Liu B.Y. Possible sarcopenia and risk of new-onset type 2 diabetes mellitus in older adults in China: A 7-year longitudinal cohort study. *BMC Geriatr.* 2023;23:404. doi: 10.1186/s12877-023-04104-9. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

13. Lee S.H., Choi H., Kim K.Y., Lee H.S., Jung J.M. Appendicular Skeletal Muscle Mass Associated with Sarcopenia as a Predictor of Poor Functional Outcomes in Ischemic Stroke. *Clin. Interv. Aging.* 2023;18:1009–1020. doi: 10.2147/CIA.S402548. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

14. Morley J.E., Anker S.D., von Haehling S. Prevalence, incidence, and clinical impact of sarcopenia: Facts, numbers, and epidemiology-update 2014. *J. Cachexia Sarcopenia Muscle*. 2014;5:253–259. doi: 10.1007/s13539-014-0161-y. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

15. Su Y., Hirayama K., Han T.F., Izutsu M., Yuki M. Sarcopenia Prevalence and Risk Factors among Japanese Community Dwelling Older Adults Living in a Snow-Covered City According to EWGSOP2. *J. Clin. Med.* 2019;8:291. doi: 10.3390/jcm8030291. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

16. Shafiee G., Keshtkar A., Soltani A., Ahadi Z., Larijani B., Heshmat R. Prevalence of sarcopenia in the world: A systematic review and meta- analysis of general population studies. *J. Diabetes Metab. Disord.* 2017;16:21. doi: 10.1186/s40200-017-0302-x. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
17. Makizako H., Nakai Y., Tomioka K., Taniguchi Y. Prevalence of sarcopenia defined using the Asia Working Group for Sarcopenia criteria in Japanese community-dwelling older adults: A systematic review and meta-analysis. *Phys. Ther. Res.* 2019;22:53–57. doi: 10.1298/ptr.R0005. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
18. Zhang X., Huang P., Dou Q., Wang C., Zhang W., Yang Y., Wang J., Xie X., Zhou J., Zeng Y. Falls among older adults with sarcopenia dwelling in nursing home or community: A meta-analysis. *Clin. Nutr.* 2020;39:33–39. doi: 10.1016/j.clnu.2019.01.002. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
19. Zhang X., Zhang W., Wang C., Tao W., Dou Q., Yang Y. Sarcopenia as a predictor of hospitalization among older people: A systematic review and meta-analysis. *BMC Geriatr.* 2018;18:188. doi: 10.1186/s12877-018-0878-0. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
20. Liu P., Hao Q., Hai S., Wang H., Cao L., Dong B. Sarcopenia as a predictor of all-cause mortality among community-dwelling older people: A systematic review and meta-analysis. *Maturitas.* 2017;103:16–22. doi: 10.1016/j.maturitas.2017.04.007. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
21. Zhang X., Wang C., Dou Q., Zhang W., Yang Y., Xie X. Sarcopenia as a predictor of all-cause mortality among older nursing home residents: A systematic review and meta-analysis. *BMJ Open.* 2018;8:e021252. doi: 10.1136/bmjopen-2017-021252. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
22. Yang M., Liu Y., Zuo Y., Tang H. Sarcopenia for predicting falls and hospitalization in community-dwelling older adults: EWGSOP versus EWGSOP2. *Sci. Rep.* 2019;9:17636. doi: 10.1038/s41598-019-53522-6. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
23. Pinedo-Villanueva R., Westbury L.D., Syddall H.E., Sanchez-Santos M.T., Dennison E.M., Robinson S.M., Cooper C. Health Care Costs Associated With Muscle Weakness: A UK Population-Based Estimate. *Calcif. Tissue Int.* 2019;104:137–144. doi: 10.1007/s00223-018-0478-1. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
24. Goates S., Du K., Arensberg M.B., Gaillard T., Guralnik J., Pereira S.L. Economic Impact of Hospitalizations in US Adults with Sarcopenia. *J. Frailty Aging.* 2019;8:93–99. doi: 10.14283/jfa.2019.10. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
25. Wall B.T., Gorissen S.H., Pennings B., Koopman R., Groen B.B., Verdijk L.B., van Loon L.J. Aging Is Accompanied by a Blunted Muscle Protein Synthetic Response to Protein Ingestion. *PLoS ONE.* 2015;10:e0140903. doi: 10.1371/journal.pone.0140903. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

26. Tan K.T., Ang S.J., Tsai S.Y. Sarcopenia: Tilting the Balance of Protein Homeostasis. *Proteomics*. 2020;20:e1800411. doi: 10.1002/pmic.201800411. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
27. Wiedmer P, Jung T, Castro J.P, Pomatto L.C.D., Sun P.Y., Davies K.J.A., Grune T. Sarcopenia—Molecular mechanisms and open questions. *Ageing Res. Rev.* 2021;65:101200. doi: 10.1016/j.arr.2020.101200. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
28. Zhang Y, Zhang P, Wang F, Xing F. Influencing factors of sarcopenia in older adults based on the Newman system model: A case-control study. *Eur. Geriatr. Med.* 2023;14:1049–1057. doi: 10.1007/s41999-023-00822-w. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
29. Robinson S., Granic A., Sayer A.A. Nutrition and Muscle Strength, As the Key Component of Sarcopenia: An Overview of Current Evidence. *Nutrients*. 2019;11:2942. doi: 10.3390/nu11122942. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
30. Calvani R., Picca A., Coelho-Júnior H.J., Tosato M., Marzetti E., Landi F. Diet for the prevention and management of sarcopenia. *Metabolism*. 2023;146:155637. doi: 10.1016/j.metabol.2023.155637. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
31. Freiburger E., Sieber C., Pfeifer K. Physical activity, exercise, and sarcopenia—Future challenges. *Wien. Med. Wochenschr.* 2011;161:416–425. doi: 10.1007/s10354-011-0001-z. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
32. Liu S., Zhang L., Li S. Advances in nutritional supplementation for sarcopenia management. *Front. Nutr.* 2023;10:1189522. doi: 10.3389/fnut.2023.1189522. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
33. Lo J.H.T., Yiu T., Ong M.T.Y., Lee W.Y.W. Sarcopenia: Current treatments and new regenerative therapeutic approaches. *J. Orthop. Transl.* 2020;23:38–52. doi: 10.1016/j.jot.2020.04.002. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
34. Landi F, Calvani R., Cesari M., Tosato M., Martone A.M., Ortolani E., Saveria G., Salini S., Sisto A., Picca A., et al. Sarcopenia: An Overview on Current Definitions, Diagnosis and Treatment. *Curr. Protein Pept. Sci.* 2018;19:633–638. doi: 10.2174/1389203718666170607113459. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
35. Deutz N.E., Bauer J.M., Barazzoni R., Biolo G., Boirie Y., Bosy-Westphal A., Cederholm T., Cruz-Jentoft A., Krznaric Z., Nair K.S., et al. Protein intake and exercise for optimal muscle function with aging: Recommendations from the ESPEN Expert Group. *Clin. Nutr.* 2014;33:929–936. doi: 10.1016/j.clnu.2014.04.007. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
36. Zeng D., Ling X.Y., Fang Z.L., Lu Y.F. Optimal exercise to improve physical ability and performance in older adults with sarcopenia: A systematic review and network meta-analysis. *Geriatr. Nurs.* 2023;52:199–207. doi: 10.1016/j.gerinurse.2023.06.005. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
37. Hurst C., Robinson S.M., Witham M.D., Dodds R.M., Granic A., Buckland C., De Biase S., Finnegan S., Rochester L., Skelton D.A., et al. Resistance exercise as a treatment for sarcopenia: Prescription and

delivery. *Age Ageing*. 2022;51:afac003. doi: 10.1093/ageing/afac003. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

38. McLeod J.C., Stokes T., Phillips S.M. Resistance Exercise Training as a Primary Countermeasure to Age-Related Chronic Disease. *Front. Physiol.* 2019;10:645. doi: 10.3389/fphys.2019.00645. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

39. Giallauria F., Cittadini A., Smart N.A., Vigorito C. Resistance training and sarcopenia. *Monaldi Arch. Chest Dis.* 2016;84:738. doi: 10.4081/monaldi.2015.738. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

40. Tessier A.J., Chevalier S. An Update on Protein, Leucine, Omega-3 Fatty Acids, and Vitamin D in the Prevention and Treatment of Sarcopenia and Functional Decline. *Nutrients*. 2018;10:1099. doi: 10.3390/nu10081099. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

41. Song Z., Pan T., Tong X., Yang Y., Zhang Z. The effects of nutritional supplementation on older sarcopenic individuals who engage in resistance training: A meta-analysis. *Front. Nutr.* 2023;10:1109789. doi: 10.3389/fnut.2023.1109789. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

42. Kuo Y.Y., Chang H.Y., Huang Y.C., Liu C.W. Effect of Whey Protein Supplementation in Postmenopausal Women: A Systematic Review and Meta-Analysis. *Nutrients*. 2022;14:4210. doi: 10.3390/nu14194210. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

43. Kirwan R.P., Mazidi M., Rodríguez García C., Lane K.E., Jafari A., Butler T., Perez de Heredia F., Davies I.G. Protein interventions augment the effect of resistance exercise on appendicular lean mass and handgrip strength in older adults: A systematic review and meta-analysis of randomized controlled trials. *Am. J. Clin. Nutr.* 2022;115:897–913. doi: 10.1093/ajcn/nqab355. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

44. Huang L.P., Condello G., Kuo C.H. Effects of Milk Protein in Resistance Training-Induced Lean Mass Gains for Older Adults Aged ≥ 60 y: A Systematic Review and Meta-Analysis. *Nutrients*. 2021;13:2815. doi: 10.3390/nu13082815. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

45. Vieira A.F., Santos J.S., Costa R.R., Cadore E.L., Macedo R.C.O. Effects of Protein Supplementation Associated with Resistance Training on Body Composition and Muscle Strength in Older Adults: A Systematic Review of Systematic Reviews with Meta-analyses. *Sports Med.* 2022;52:2511–2522. doi: 10.1007/s40279-022-01704-0. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

46. Park S.H., Roh Y. Which intervention is more effective in improving sarcopenia in older adults? A systematic review with meta-analysis of randomized controlled trials. *Mech. Ageing Dev.* 2023;210:111773. doi: 10.1016/j.mad.2022.111773. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

47. Carcelén-Fraile M.D.C., Lorenzo-Nocino M.F., Afanador-Restrepo D.F., Rodríguez-López C., Aibar-Almazán A., Hita-Contreras F., Achalandabaso-Ochoa A., Castellote-Caballero Y. Effects of different intervention combined with resistance training on musculoskeletal health in older male adults with

sarcopenia: A systematic review. *Front. Public Health.* 2023;10:1037464. doi:

10.3389/fpubh.2022.1037464. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

48. Negm A.M., Lee J., Hamidian R., Jones C.A., Khadaroo R.G. Management of Sarcopenia: A Network Meta-Analysis of Randomized Controlled Trials. *J. Am. Med. Dir. Assoc.* 2022;23:707–714. doi: 10.1016/j.jamda.2022.01.057. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

49. Liao C.D., Wu Y.T., Tsao J.Y., Chen P.R., Tu Y.K., Chen H.C., Liou T.H. Effects of Protein Supplementation Combined with Exercise Training on Muscle Mass and Function in Older Adults with Lower-Extremity Osteoarthritis: A Systematic Review and Meta-Analysis of Randomized Trials. *Nutrients.* 2020;12:2422. doi: 10.3390/nu12082422. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

50. Cuyul-Vásquez I., Pezo-Navarrete J., Vargas-Arriagada C., Ortega-Díaz C., Sepúlveda-Loyola W., Hirabara S.M., Marzuca-Nassr G.N. Effectiveness of Whey Protein Supplementation during Resistance Exercise Training on Skeletal Muscle Mass and Strength in Older People with Sarcopenia: A Systematic Review and Meta-Analysis. *Nutrients.* 2023;15:3424. doi: 10.3390/nu15153424. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

51. Liao C.D., Tsao J.Y., Wu Y.T., Cheng C.P., Chen H.C., Huang Y.C., Chen H.C., Liou T.H. Effects of protein supplementation combined with resistance exercise on body composition and physical function in older adults: A systematic review and meta-analysis. *Am. J. Clin. Nutr.* 2017;106:1078–1091. doi: 10.3945/ajcn.116.143594. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

52. Murphy C.H., Saddler N.I., Devries M.C., McGlory C., Baker S.K., Phillips S.M. Leucine supplementation enhances integrative myofibrillar protein synthesis in free-living older men consuming lower- and higher-protein diets: A parallel-group crossover study. *Am. J. Clin. Nutr.* 2016;104:1594–1606. doi: 10.3945/ajcn.116.136424. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

53. Trommelen J., Betz M.W., van Loon L.J.C. The Muscle Protein Synthetic Response to Meal Ingestion Following Resistance-Type Exercise. *Sports Med.* 2019;49:185–197. doi: 10.1007/s40279-019-01053-5. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

54. Landi F., Calvani R., Tosato M., Martone A.M., Ortolani E., Saveria G., D'Angelo E., Sisto A., Marzetti E. Protein Intake and Muscle Health in Old Age: From Biological Plausibility to Clinical Evidence. *Nutrients.* 2016;8:295. doi: 10.3390/nu8050295. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

55. Reitelseder S., Dideriksen K., Agergaard J., Malmgaard-Clausen N.M., Bechshoeft R.L., Petersen R.K., Serena A., Mikkelsen U.R., Holm L. Even effect of milk protein and carbohydrate intake but no further effect of heavy resistance exercise on myofibrillar protein synthesis in older men. *Eur. J. Nutr.* 2019;58:583–595. doi: 10.1007/s00394-018-1641-1. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

56. Pinckaers P.J.M., Kouw I.W.K., Gorissen S.H.M., Houben L.H.P., Senden J.M., Wodzig W., de Groot L., Verdijk L.B., Snijders T., van Loon L.J.C. The Muscle Protein Synthetic Response to the Ingestion of a Plant-Derived Protein Blend Does Not Differ from an Equivalent Amount of Milk Protein in Healthy

Young Males. *J. Nutr.* 2023;152:2734–2743. doi: 10.1093/jn/nxac222. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

57. Lanng S.K., Oxfeldt M., Pedersen S.S., Johansen F.T., Risikesan J., Lejel T., Bertram H.C., Hansen M. Influence of protein source (cricket, pea, whey) on amino acid bioavailability and activation of the mTORC1 signaling pathway after resistance exercise in healthy young males. *Eur. J. Nutr.* 2023;62:1295–1308. doi: 10.1007/s00394-022-03071-y. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

58. Colonetti T., Grande A.J., Milton K., Foster C., Alexandre M.C., Uggioni M.L., Rosa M.I. Effects of whey protein supplement in the elderly submitted to resistance training: Systematic review and meta-analysis. *Int. J. Food Sci. Nutr.* 2017;68:257–264. doi: 10.1080/09637486.2016.1232702. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

59. Monteyne A.J., Coelho M.O.C., Porter C., Abdelrahman D.R., Jameson T.S.O., Jackman S.R., Blackwell J.R., Finnigan T.J.A., Stephens F.B., Dirks M.L., et al. Mycoprotein ingestion stimulates protein synthesis rates to a greater extent than milk protein in rested and exercised skeletal muscle of healthy young men: A randomized controlled trial. *Am. J. Clin. Nutr.* 2020;112:318–333. doi: 10.1093/ajcn/nqaa092. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

60. West S., Monteyne A.J., Whelehan G., van der Heijden I., Abdelrahman D.R., Murton A.J., Finnigan T.J.A., Stephens F.B., Wall B.T. Ingestion of mycoprotein, pea protein, and their blend support comparable postexercise myofibrillar protein synthesis rates in resistance-trained individuals. *Am. J. Physiol. Endocrinol. Metab.* 2023;325:E267–E279. doi: 10.1152/ajpendo.00166.2023. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

61. Burd N.A., Yang Y., Moore D.R., Tang J.E., Tarnopolsky M.A., Phillips S.M. Greater stimulation of myofibrillar protein synthesis with ingestion of whey protein isolate v. micellar casein at rest and after resistance exercise in elderly men. *Br. J. Nutr.* 2012;108:958–962. doi: 10.1017/S0007114511006271. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

62. Miller P.E., Alexander D.D., Perez V. Effects of Whey Protein and Resistance Exercise on Body Composition: A Meta-Analysis of Randomized Controlled Trials. *J. Am. Coll. Nutr.* 2014;33:163–175. doi: 10.1080/07315724.2013.875365. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

63. Messina M., Lynch H., Dickinson J.M., Reed K.E. No Difference Between the Effects of Supplementing With Soy Protein Versus Animal Protein on Gains in Muscle Mass and Strength in Response to Resistance Exercise. *Int. J. Sport Nutr. Exerc. Metab.* 2018;28:674–685. doi: 10.1123/ijsnem.2018-0071. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

64. Valenzuela P.L., Mata F., Morales J.S., Castillo-García A., Lucia A. Does Beef Protein Supplementation Improve Body Composition and Exercise Performance? A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Nutrients.* 2019;11:1429. doi: 10.3390/nu11061429. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

65. Zanini B., Simonetto A., Zubani M., Castellano M., Gilioli G. The Effects of Cow-Milk Protein Supplementation in Elderly Population: Systematic Review and Narrative Synthesis. *Nutrients.*

2020;12:2548. doi: 10.3390/nu12092548. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

66. Piri Damaghi M., Mirzababaei A., Moradi S., Daneshzad E., Tavakoli A., Clark C.C.T., Mirzaei K. Comparison of the effect of soya protein and whey protein on body composition: A meta-analysis of randomised clinical trials. *Br. J. Nutr.* 2022;127:885–895. doi: 10.1017/S0007114521001550. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

67. Morgan P.T., Harris D.O., Marshall R.N., Quinlan J.I., Edwards S.J., Allen S.L., Breen L. Protein Source and Quality for Skeletal Muscle Anabolism in Young and Older Adults: A Systematic Review and Meta-Analysis. *J. Nutr.* 2021;151:1901–1920. doi: 10.1093/jn/nxab055. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

68. Shamseer L., Moher D., Clarke M., Gherzi D., Liberati A., Petticrew M., Shekelle P., Stewart L.A. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: Elaboration and explanation. *BMJ.* 2015;349:g7647. doi: 10.1136/bmj.g7647. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

69. Hutton B., Salanti G., Caldwell D.M., Chaimani A., Schmid C.H., Cameron C., Ioannidis J.P., Straus S., Thorlund K., Jansen J.P., et al. The PRISMA extension statement for reporting of systematic reviews incorporating network meta-analyses of health care interventions: Checklist and explanations. *Ann. Intern. Med.* 2015;162:777–784. doi: 10.7326/M14-2385. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

70. Hutton B., Catalá-López F., Moher D. The PRISMA statement extension for systematic reviews incorporating network meta-analysis: PRISMA-NMA. *Med. Clin.* 2016;147:262–266. doi: 10.1016/j.medcli.2016.02.025. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

71. Higgins J.P.T., Thomas J., Chandler J., Cumpston M., Li T., Page M.J., Welch V.A. *Cochrane Handbook for Systematic Reviews of Interventions Version 6.4 (Updated August 2023)*, version 6.4 (updated August 2023) ed. The Cochrane Collaboration; London, UK: 2023. [[Google Scholar](#)]

72. Higgins J.P., Altman D.G., Gøtzsche P.C., Jüni P., Moher D., Oxman A.D., Savovic J., Schulz K.F., Weeks L., Sterne J.A. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ.* 2011;343:d5928. doi: 10.1136/bmj.d5928. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

73. Begg C.B., Mazumdar M. Operating characteristics of a rank correlation test for publication bias. *Biometrics.* 1994;50:1088–1101. doi: 10.2307/2533446. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

74. Higgins J.P.T., Li T., Deeks J.J. Chapter 6: Choosing effect measures and computing estimates of effect. In: Higgins J.P.T., Thomas J., Chandler J., Cumpston M., Li T., Page M.J., Welch V.A., editors. *Cochrane Handbook for Systematic Reviews of Interventions Version 6.4 (Updated August 2023)* Cochrane; London, UK: 2023. [[Google Scholar](#)]

75. Wan X., Wang W., Liu J., Tong T. Estimating the sample mean and standard deviation from the sample size, median, range and/or interquartile range. *BMC Med. Res. Methodol.* 2014;14:135. doi: 10.1186/1471-2288-14-135. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

76. Cohen J. Statistical Power Analysis for the Behavioral Sciences. 2nd ed. Lawrence Erlbaum Associates; Hillsdale, MI, USA: 1988. [[Google Scholar](#)]
77. Chaimani A., Caldwell D.M., Li T., Higgins J.P.T.S.G. Chapter 11: Undertaking network meta-analyses. In: Higgins J.P.T., Thomas J., Chandler J., Cumpston M., Li T., Page M.J., Welch V.A., editors. Cochrane Handbook for Systematic Reviews of Interventions Version 6.3 (Updated February 2022) Cochrane; London, UK: 2022. [[Google Scholar](#)]
78. Jackson D., Boddington P., White I.R. The design-by-treatment interaction model: A unifying framework for modelling loop inconsistency in network meta-analysis. Res. Synth. Methods. 2016;7:329–332. doi: 10.1002/jrsm.1188. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
79. Salanti G., Ades A.E., Ioannidis J.P. Graphical methods and numerical summaries for presenting results from multiple-treatment meta-analysis: An overview and tutorial. J. Clin. Epidemiol. 2011;64:163–171. doi: 10.1016/j.jclinepi.2010.03.016. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
80. Harrer M., Cuijpers P., Furukawa T.A., Ebert D.D. Chapter 12: Network Meta-Analysis. In: Harrer M., Cuijpers P., Furukawa T.A., Ebert D.D., editors. Doing Meta-Analysis with R: A Hands-On Guide. Chapman & Hall; London, UK: CRC Press; Boca Raton, FL, USA: 2021. [[Google Scholar](#)]
81. Li G., Xie F., Yan S., Hu X., Jin B., Wang J., Wu J., Yin D., Xie Q. Subhealth: Definition, criteria for diagnosis and potential prevalence in the central region of China. BMC Public Health. 2013;13:446. doi: 10.1186/1471-2458-13-446. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
82. Shim S.R., Kim S.-J., Lee J., Rücker G. Network meta-analysis: Application and practice using R software. Epidemiol. Health. 2019;41:e2019013. doi: 10.4178/epih.e2019013. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
83. Brignardello-Petersen R., Bonner A., Alexander P.E., Siemieniuk R.A., Furukawa T.A., Rochwerg B., Hazlewood G.S., Alhazzani W., Mustafa R.A., Murad M.H., et al. Advances in the GRADE approach to rate the certainty in estimates from a network meta-analysis. J. Clin. Epidemiol. 2018;93:36–44. doi: 10.1016/j.jclinepi.2017.10.005. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
84. Papakonstantinou T., Nikolakopoulou A., Rücker G., Chaimani A., Schwarzer G., Egger M., Salanti G. Estimating the contribution of studies in network meta-analysis: Paths, flows and streams. F1000Res. 2018;7:610. doi: 10.12688/f1000research.14770.1. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
85. Furukawa T.A., Miura T., Chaimani A., Leucht S., Cipriani A., Noma H., Mitsuyasu H., Kanba S., Salanti G. Using the contribution matrix to evaluate complex study limitations in a network meta-analysis: A case study of bipolar maintenance pharmacotherapy review. BMC Res. Notes. 2016;9:218. doi: 10.1186/s13104-016-2019-1. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
86. Salanti G., Del Giovane C., Chaimani A., Caldwell D.M., Higgins J.P. Evaluating the quality of evidence from a network meta-analysis. PLoS ONE. 2014;9:e99682. doi: 10.1371/journal.pone.0099682. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

87. Chiocchia V, Nikolakopoulou A, Higgins J.P.T., Page M.J., Papakonstantinou T, Cipriani A, Furukawa T.A., Siontis G.C.M., Egger M., Salanti G. ROB-MEN: A tool to assess risk of bias due to missing evidence in network meta-analysis. *BMC Med.* 2021;19:304. doi: 10.1186/s12916-021-02166-3. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
88. Aas S.N., Seynnes O., Benestad H.B., Raastad T. Strength training and protein supplementation improve muscle mass, strength, and function in mobility-limited older adults: A randomized controlled trial. *Aging Clin. Exp. Res.* 2019;32:605–616. doi: 10.1007/s40520-019-01234-2. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
89. Amasene M., Besga A., Echeverria I., Urquiza M., Ruiz J.R., Rodriguez-Larrad A., Aldamiz M., Anaut P., Irazusta J., Labayen I. Effects of Leucine-Enriched Whey Protein Supplementation on Physical Function in Post-Hospitalized Older Adults Participating in 12-Weeks of Resistance Training Program: A Randomized Controlled Trial. *Nutrients.* 2019;11:2337. doi: 10.3390/nu11102337. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
90. Amasene M., Cadenas-Sanchez C., Echeverria I., Sanz B., Alonso C., Tobalina I., Irazusta J., Labayen I., Besga A. Effects of Resistance Training Intervention along with Leucine-Enriched Whey Protein Supplementation on Sarcopenia and Frailty in Post-Hospitalized Older Adults: Preliminary Findings of a Randomized Controlled Trial. *J. Clin. Med.* 2021;11:97. doi: 10.3390/jcm11010097. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
91. Arciero P.J., Baur D., Connelly S., Ormsbee M.J. Timed-daily ingestion of whey protein and exercise training reduces visceral adipose tissue mass and improves insulin resistance: The PRISE study. *J. Appl. Physiol.* 2014;117:1–10. doi: 10.1152/jappphysiol.00152.2014. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
92. Arnarson A., Gudny Geirsdottir O., Ramel A., Briem K., Jonsson P.V., Thorsdottir I. Effects of whey proteins and carbohydrates on the efficacy of resistance training in elderly people: Double blind, randomised controlled trial. *Eur. J. Clin. Nutr.* 2013;67:821–826. doi: 10.1038/ejcn.2013.40. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
93. Bagheri R., Hooshmand Moghadam B., Candow D.G., Elliott B.T., Wong A., Ashtary-Larky D., Forbes S.C., Rashidlamir A. Effects of Icelandic yogurt consumption and resistance training in healthy untrained older males. *Br. J. Nutr.* 2022;127:1334–1342. doi: 10.1017/S0007114521002166. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
94. Bembien M.G., Witten M.S., Carter J.M., Eliot K.A., Knehans A.W., Bembien D.A. The effects of supplementation with creatine and protein on muscle strength following a traditional resistance training program in middle-aged and older men. *J. Nutr. Health Aging.* 2010;14:155–159. doi: 10.1007/s12603-009-0124-8. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
95. Bijeh N., Mohammadnia-Ahmadi M., Hooshmand-Moghadam B., Eskandari M., Golestani F. Effects of Soy Milk in Conjunction With Resistance Training on Physical Performance and Skeletal Muscle

Regulatory Markers in Older Men. *Biol. Res. Nurs.* 2022;24:294–307. doi: 10.1177/10998004211073123. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

96. Candow D.G., Chilibeck P.D., Facci M., Abeysekara S., Zello G.A. Protein supplementation before and after resistance training in older men. *Eur. J. Appl. Physiol.* 2006;97:548–556. doi: 10.1007/s00421-006-0223-8. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

97. Candow D.G., Little J.P., Chilibeck P.D., Abeysekara S., Zello G.A., Kazachkov M., Cornish S.M., Yu P.H. Low-dose creatine combined with protein during resistance training in older men. *Med. Sci. Sports Exerc.* 2008;40:1645–1652. doi: 10.1249/MSS.0b013e318176b310. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

98. Chale A., Cloutier G.J., Hau C., Phillips E.M., Dallal G.E., Fielding R.A. Efficacy of whey protein supplementation on resistance exercise-induced changes in lean mass, muscle strength, and physical function in mobility-limited older adults. *J. Gerontol. Ser. A Biol. Sci. Med. Sci.* 2013;68:682–690. doi: 10.1093/gerona/gls221. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

99. Chang Y.M. Effects of resistance training combined with nutritional intervention on alleviating sarcopenia in the elderly. *Electron. J. Pract. Clin. Nurs. Sci.* 2019;4:48–50. [[Google Scholar](#)]

100. Chen S., Sheng Y.L., Qi T., Sheng X.M., Jin X.W. Effects of fortified nutrition combining with resistance exercise on physical function and activity of daily living for elderly patients with sarcopenia. *J. Nurs. Sci.* 2017;32:8–10. doi: 10.3870/j.issn.1001-4152.2017.21.008. [[DOI](#)] [[Google Scholar](#)]

101. Colonetti T., Grande A.J., da Rocha F.R., Ronconi Dondossola E., Tuon L., Gomes Batista Teles H., Minotto Bom B., Colonetti L., da Rosa M.I. Whey protein and vitamin D supplementation in institutionalized older adults: A randomized trial. *Nutr. Health.* 2023;29:129–138. doi: 10.1177/02601060211060665. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

102. Daly R.M., O’Connell S.L., Mundell N.L., Grimes C.A., Dunstan D.W., Nowson C.A. Protein-enriched diet, with the use of lean red meat, combined with progressive resistance training enhances lean tissue mass and muscle strength and reduces circulating IL-6 concentrations in elderly women: A cluster randomized controlled trial. *Am. J. Clin. Nutr.* 2014;99:899–910. doi: 10.3945/ajcn.113.064154. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

103. de Azevedo Bach S., Radaelli R., Beck Schemes M., Neske R., Garbelotto C., Roschel H., Silveira Pinto R., Dornelles Schneider C. Can supplemental protein to low-protein containing meals superimpose on resistance-training muscle adaptations in older adults? A randomized clinical trial. *Exp. Gerontol.* 2022;162:111760. doi: 10.1016/j.exger.2022.111760. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

104. de Carvalho Bastone A., Nobre L.N., de Souza Moreira B., Rosa I.F., Ferreira G.B., Santos D.D.L., Monteiro N., Alves M.D., Gandra R.A., de Lira E.M. Independent and combined effect of home-based progressive resistance training and nutritional supplementation on muscle strength, muscle mass and physical function in dynapenic older adults with low protein intake: A randomized controlled trial.

Arch. Gerontol. Geriatr. 2020;89:104098. doi: 10.1016/j.archger.2020.104098. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

105. Deer R.R., Dickinson J.M., Baillargeon J., Fisher S.R., Raji M., Volpi E. A Phase I Randomized Clinical Trial of Evidence-Based, Pragmatic Interventions to Improve Functional Recovery After Hospitalization in Geriatric Patients. *J. Gerontol. Ser. A Biol. Sci. Med. Sci.* 2019;74:1628–1636. doi: 10.1093/gerona/glz084. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

106. Deibert P., Solleder F., König D., Vitolins M.Z., Dickhuth H.H., Gollhofer A., Berg A. Soy protein based supplementation supports metabolic effects of resistance training in previously untrained middle aged males. *Aging Male Off. J. Int. Soc. Study Aging Male.* 2011;14:273–279. doi: 10.3109/13685538.2011.565091. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

107. Dirks M.L., Tieland M., Verdijk L.B., Losen M., Nilwik R., Mensink M., de Groot L.C.P.G.M., van Loon L.J.C. Protein Supplementation Augments Muscle Fiber Hypertrophy but Does Not Modulate Satellite Cell Content During Prolonged Resistance-Type Exercise Training in Frail Elderly. *J. Am. Med. Dir. Assoc.* 2017;18:608–615. doi: 10.1016/j.jamda.2017.02.006. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

108. Duff W.R., Chilibeck P.D., Rooke J.J., Kaviani M., Krentz J.R., Haines D.M. The effect of bovine colostrum supplementation in older adults during resistance training. *Int. J. Sport Nutr. Exerc. Metab.* 2014;24:276–285. doi: 10.1123/ijsnem.2013-0182. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

109. Dulac M.C., Pion C.H., Lemieux F.C., Pinheiro Carvalho L., El Hajj Boutros G., Bélanger M., Gaudreau P., Chevalier S., Morais J.A., Noirez P., et al. Effects of slow- v. fast-digested protein supplementation combined with mixed power training on muscle function and functional capacities in older men. *Br. J. Nutr.* 2021;125:1017–1033. doi: 10.1017/S0007114520001932. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

110. Eliot K.A., Knehans A.W., Bembien D.A., Witten M.S., Carter J., Bembien M.G. The effects of creatine and whey protein supplementation on body composition in men aged 48 to 72 years during resistance training. *J. Nutr. Health Aging.* 2008;12:208–212. doi: 10.1007/BF02982622. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

111. Fiatarone M.A., O'Neill E.F., Ryan N.D., Clements K.M., Solares G.R., Nelson M.E., Roberts S.B., Kehayias J.J., Lipsitz L.A., Evans W.J. Exercise Training and Nutritional Supplementation for Physical Frailty in Very Elderly People. *N. Engl. J. Med.* 1994;330:1769–1775. doi: 10.1056/NEJM199406233302501. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

112. Formica M.B., Gianoudis J., Nowson C.A., O'Connell S.L., Milte C., Ellis K.A., Daly R.M. Effect of lean red meat combined with a multicomponent exercise program on muscle and cognitive function in older adults: A 6-month randomized controlled trial. *Am. J. Clin. Nutr.* 2020;112:113–128. doi: 10.1093/ajcn/nqaa104. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

113. Francis P., Mc Cormack W., Toomey C., Norton C., Saunders J., Kerin E., Lyons M., Jakeman P. Twelve weeks' progressive resistance training combined with protein supplementation beyond habitual

intakes increases upper leg lean tissue mass, muscle strength and extended gait speed in healthy older women. *Biogerontology*. 2017;18:881–891. doi: 10.1007/s10522-016-9671-7. [[DOI ↗](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar ↗](#)]

114. Gade J., Beck A.M., Andersen H.E., Christensen B., Rønholt F., Klausen T.W., Vinther A., Astrup A. Protein supplementation combined with low-intensity resistance training in geriatric medical patients during and after hospitalisation: A randomised, double-blind, multicentre trial. *Br. J. Nutr.* 2019;122:1006–1020. doi: 10.1017/S0007114519001831. [[DOI ↗](#)] [[PubMed](#)] [[Google Scholar ↗](#)]

115. Granic A., Hurst C., Dismore L., Stevenson E., Sayer A.A., Aspray T. Feasibility and acceptability of a milk and resistance exercise intervention to improve muscle function in community-dwelling older adults (MilkMAN): Pilot study. *PLoS ONE*. 2020;15:e0235952. doi: 10.1371/journal.pone.0235952. [[DOI ↗](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar ↗](#)]

116. Griffen C., Duncan M., Hattersley J., Weickert M.O., Dallaway A., Renshaw D. Effects of resistance exercise and whey protein supplementation on skeletal muscle strength, mass, physical function, and hormonal and inflammatory biomarkers in healthy active older men: A randomised, double-blind, placebo-controlled trial. *Exp. Gerontol.* 2022;158:111651. doi: 10.1016/j.exger.2021.111651. [[DOI ↗](#)] [[PubMed](#)] [[Google Scholar ↗](#)]

117. Gronstedt H., Vikstrom S., Cederholm T., Franzen E., Luiking Y.C., Seiger A., Wimo A., Faxen-Ingving G., Bostrom A.M. Effect of Sit-to-Stand Exercises Combined With Protein-Rich Oral Supplementation in Older Persons: The Older Person's Exercise and Nutrition Study. *J. Am. Med. Dir. Assoc.* 2020;21:1229–1237. doi: 10.1016/j.jamda.2020.03.030. [[DOI ↗](#)] [[PubMed](#)] [[Google Scholar ↗](#)]

118. Hamarsland H., Johansen M.K., Seeberg F., Brochmann M., Garthe I., Benestad H.B., Raastad T. Native Whey Induces Similar Adaptation to Strength Training as Milk, despite Higher Levels of Leucine, in Elderly Individuals. *Nutrients*. 2019;11:2094. doi: 10.3390/nu11092094. [[DOI ↗](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar ↗](#)]

119. Haß U., Kochlik B., Herpich C., Rudloff S., Norman K. Effects of an Omega-3 Supplemented, High-Protein Diet in Combination with Vibration and Resistance Exercise on Muscle Power and Inflammation in Old Adults: A Pilot Randomized Controlled Trial. *Nutrients*. 2022;14:4274. doi: 10.3390/nu14204274. [[DOI ↗](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar ↗](#)]

120. Haub M.D., Wells A.M., Campbell W.W. Beef and soy-based food supplements differentially affect serum lipoprotein-lipid profiles because of changes in carbohydrate intake and novel nutrient intake ratios in older men who resistive-train. *Metabolism*. 2005;54:769–774. doi: 10.1016/j.metabol.2005.01.019. [[DOI ↗](#)] [[PubMed](#)] [[Google Scholar ↗](#)]

121. Haub M.D., Wells A.M., Tarnopolsky M.A., Campbell W.W. Effect of protein source on resistive-training-induced changes in body composition and muscle size in older men. *Am. J. Clin. Nutr.* 2002;76:511–517. doi: 10.1093/ajcn/76.3.511. [[DOI ↗](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar ↗](#)]

122. He L.N., Li D.M. Observation on the Clinical Effect of Dietary Nutrition Intervention Combined with Anti-resistance Training in Elderly Male Patients with Sarcopenia. *Smart Healthc.* 2022;8:44–47. doi: 10.19335/j.cnki.2096-1219.2022.19.011. [[DOI ↗](#)] [[Google Scholar ↗](#)]
123. Holm L., Olesen J.L., Matsumoto K., Doi T., Mizuno M., Alsted T.J., Mackey A.L., Schwarz P., Kjaer M. Protein-containing nutrient supplementation following strength training enhances the effect on muscle mass, strength, and bone formation in postmenopausal women. *J. Appl. Physiol.* 2008;105:274–281. doi: 10.1152/jappphysiol.00935.2007. [[DOI ↗](#)] [[PubMed](#)] [[Google Scholar ↗](#)]
124. Holwerda A.M., Overkamp M., Paulussen K.J.M., Smeets J.S.J., van Kranenburg J., Backx E.M.P., Gijzen A.P., Goessens J.P.B., Verdijk L.B., van Loon L.J.C. Protein Supplementation after Exercise and before Sleep Does Not Further Augment Muscle Mass and Strength Gains during Resistance Exercise Training in Active Older Men. *J. Nutr.* 2018;148:1723–1732. doi: 10.1093/jn/nxy169. [[DOI ↗](#)] [[PubMed](#)] [[Google Scholar ↗](#)]
125. Kang Y., Kim N., Choi Y.J., Lee Y., Yun J., Park S.J., Park H.S., Chung Y.S., Park Y.K. Leucine-Enriched Protein Supplementation Increases Lean Body Mass in Healthy Korean Adults Aged 50 Years and Older: A Randomized, Double-Blind, Placebo-Controlled Trial. *Nutrients.* 2020;12:1816. doi: 10.3390/nu12061816. [[DOI ↗](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar ↗](#)]
126. Karelis A.D., Messier V., Suppere C., Briand P., Rabasa-Lhoret R. Effect of cysteine-rich whey protein (immunocal(R)) supplementation in combination with resistance training on muscle strength and lean body mass in non-frail elderly subjects: A randomized, double-blind controlled study. *J. Nutr. Health Aging.* 2015;19:531–536. doi: 10.1007/s12603-015-0442-y. [[DOI ↗](#)] [[PubMed](#)] [[Google Scholar ↗](#)]
127. Kemmler W., Kohl M., Fröhlich M., Jakob F., Engelke K., von Stengel S., Schoene D. Effects of High-Intensity Resistance Training on Osteopenia and Sarcopenia Parameters in Older Men with Osteosarcopenia-One-Year Results of the Randomized Controlled Franconian Osteopenia and Sarcopenia Trial (FrOST) *J Bone Min. Res.* 2020;35:1634–1644. doi: 10.1002/jbmr.4027. [[DOI ↗](#)] [[PubMed](#)] [[Google Scholar ↗](#)]
128. Kemmler W., Kohl M., Jakob F., Engelke K., von Stengel S. Effects of High Intensity Dynamic Resistance Exercise and Whey Protein Supplements on Osteosarcopenia in Older Men with Low Bone and Muscle Mass. Final Results of the Randomized Controlled FrOST Study. *Nutrients.* 2020;12:234. doi: 10.3390/nu12082341. [[DOI ↗](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar ↗](#)]
129. Krause M., Crognale D., Cogan K.E., Contarelli S., Egan B., Newsholme P., De Vito G. The effects of a combined bodyweight-based and elastic bands resistance training, with or without protein supplementation, on muscle mass, signaling and heat shock response in healthy older people. *Exp. Gerontol.* 2019;115:104–113. doi: 10.1016/j.exger.2018.12.004. [[DOI ↗](#)] [[PubMed](#)] [[Google Scholar ↗](#)]
130. Kukuljan S., Nowson C.A., Bass S.L., Sanders K., Nicholson G.C., Seibel M.J., Salmon J., Daly R.M. Effects of a multi-component exercise program and calcium-vitamin-D3-fortified milk on bone mineral

density in older men: A randomised controlled trial. *Osteoporos. Int.* 2009;20:1241–1251. doi: 10.1007/s00198-008-0776-y. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

131. Kukuljan S., Nowson C.A., Sanders K., Daly R.M. Effects of resistance exercise and fortified milk on skeletal muscle mass, muscle size, and functional performance in middle-aged and older men: An 18-mo randomized controlled trial. *J. Appl. Physiol.* 2009;107:1864–1873. doi: 10.1152/jappphysiol.00392.2009. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

132. Kwon J., Yoshida Y., Yoshida H., Kim H., Suzuki T., Lee Y. Effects of a combined physical training and nutrition intervention on physical performance and health-related quality of life in prefrail older women living in the community: A randomized controlled trial. *J. Am. Med. Dir. Assoc.* 2015;16:e261–e268. doi: 10.1016/j.jamda.2014.12.005. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

133. Lamb D.A., Moore J.H., Smith M.A., Vann C.G., Osburn S.C., Rupple B.A., Fox C.D., Smith K.S., Altonji O.M., Power Z.M., et al. The effects of resistance training with or without peanut protein supplementation on skeletal muscle and strength adaptations in older individuals. *J. Int. Soc. Sports Nutr.* 2020;17:66. doi: 10.1186/s12970-020-00397-y. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

134. Leenders M., Verdijk L.B., Van der Hoeven L., Van Kranenburg J., Nilwik R., Wodzig W.K., Senden J.M., Keizer H.A., Van Loon L.J. Protein supplementation during resistance-type exercise training in the elderly. *Med. Sci. Sports Exerc.* 2013;45:542–552. doi: 10.1249/MSS.0b013e318272fcdb. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

135. Liao C.D., Liao Y.H., Liou T.H., Hsieh C.Y., Kuo Y.C., Chen H.C. Effects of Protein-Rich Nutritional Composition Supplementation on Sarcopenia Indices and Physical Activity during Resistance Exercise Training in Older Women with Knee Osteoarthritis. *Nutrients.* 2021;13:2487. doi: 10.3390/nu13082487. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

136. Maesta N., Nahas E.A., Nahas-Neto J., Orsatti F.L., Fernandes C.E., Traiman P., Burini R.C. Effects of soy protein and resistance exercise on body composition and blood lipids in postmenopausal women. *Maturitas.* 2007;56:350–358. doi: 10.1016/j.maturitas.2006.10.001. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

137. Maltais M.L., Ladouceur J.P., Dionne I.J. The Effect of Resistance Training and Different Sources of Postexercise Protein Supplementation on Muscle Mass and Physical Capacity in Sarcopenic Elderly Men. *J. Strength Cond. Res./Natl. Strength Cond. Assoc.* 2016;30:1680–1687. doi: 10.1519/JSC.0000000000001255. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

138. McKenna C.F., Salvador A.F., Hughes R.L., Scaroni S.E., Alamilla R.A., Askow A.T., Paluska S.A., Dilger A.C., Holscher H.D., De Lisio M. Higher protein intake during resistance training does not potentiate strength, but modulates gut microbiota, in middle-aged adults: A randomized control trial. *Am J Physiol.-Endocrinol. Metab.* 2021;320:E900–E913. doi: 10.1152/ajpendo.00574.2020. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

139. Mertz K.H., Reitelseder S., Bechshoeft R., Bulow J., Højfeldt G., Jensen M., Schacht S.R., Lind M.V., Rasmussen M.A., Mikkelsen U.R., et al. The effect of daily protein supplementation, with or without resistance training for 1 year, on muscle size, strength, and function in healthy older adults: A randomized controlled trial. *Am. J. Clin. Nutr.* 2021;113:790–800. doi: 10.1093/ajcn/nqaa372. [[DOI ↗](#)] [[PubMed](#)] [[Google Scholar ↗](#)]
140. Miller E.G., Nowson C.A., Dunstan D.W., Kerr D.A., Menzies D., Daly R.M. Effects of whey protein plus vitamin D supplementation combined with progressive resistance training on glycaemic control, body composition, muscle function and cardiometabolic risk factors in middle-aged and older overweight/obese adults with type 2 diabetes: A 24-week randomized controlled trial. *Diabetes Obes. Metab.* 2021;23:938–949. doi: 10.1111/dom.14299. [[DOI ↗](#)] [[PubMed](#)] [[Google Scholar ↗](#)]
141. Mitchell C.J., D’Souza R.F., Mitchell S.M., Figueiredo V.C., Miller B.F., Hamilton K.L., Peelor F.F., 3rd, Coronet M., Pileggi C.A., Durainayagam B., et al. Impact of dairy protein during limb immobilization and recovery on muscle size and protein synthesis; a randomized controlled trial. *J. Appl. Physiol.* 2018;124:717–728. doi: 10.1152/jappphysiol.00803.2017. [[DOI ↗](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar ↗](#)]
142. Molnar A., Jonasne Sztruhar I., Csontos A.A., Ferencz C., Varbiro S., Szekacs B. Special nutrition intervention is required for muscle protective efficacy of physical exercise in elderly people at highest risk of sarcopenia. *Physiol. Int.* 2016;103:368–376. doi: 10.1556/2060.103.2016.3.12. [[DOI ↗](#)] [[PubMed](#)] [[Google Scholar ↗](#)]
143. Mori H., Tokuda Y. Effect of whey protein supplementation after resistance exercise on the muscle mass and physical function of healthy older women: A randomized controlled trial. *Geriatr. Gerontol. Int.* 2018;18:1398–1404. doi: 10.1111/ggi.13499. [[DOI ↗](#)] [[PubMed](#)] [[Google Scholar ↗](#)]
144. Mori H., Tokuda Y. De-Training Effects Following Leucine-Enriched Whey Protein Supplementation and Resistance Training in Older Adults with Sarcopenia: A Randomized Controlled Trial with 24 Weeks of Follow-Up. *J. Nutr. Health Aging.* 2022;26:994–1002. doi: 10.1007/s12603-022-1853-1. [[DOI ↗](#)] [[PubMed](#)] [[Google Scholar ↗](#)]
145. Nabuco H.C.G., Tomeleri C.M., Fernandes R.R., Sugihara Junior P, Cavalcante E.F, Cunha P.M., Antunes M., Nunes J.P, Venturini D., Barbosa D.S., et al. Effect of whey protein supplementation combined with resistance training on body composition, muscular strength, functional capacity, and plasma-metabolism biomarkers in older women with sarcopenic obesity: A randomized, double-blind, placebo-controlled trial. *Clin. Nutr. ESPEN.* 2019;32:88–95. doi: 10.1016/j.clnesp.2019.04.007. [[DOI ↗](#)] [[PubMed](#)] [[Google Scholar ↗](#)]
146. Nabuco H.C.G., Tomeleri C.M., Fernandes R.R., Sugihara Junior P, Cavalcante E.F, Venturini D., Barbosa D.S., Silva A.M., Sardinha L.B., Cyrino E.S. Effects of Protein Intake Beyond Habitual Intakes Associated With Resistance Training on Metabolic Syndrome-Related Parameters, Isokinetic Strength, and Body Composition in Older Women. *J. Aging Phys. Act.* 2019;27:545–552. doi: 10.1123/japa.2018-0370. [[DOI ↗](#)] [[PubMed](#)] [[Google Scholar ↗](#)]

147. Nabuco H.C.G., Tomeleri C.M., Sugihara Junior P, Fernandes R.R., Cavalcante E.F., Antunes M., Ribeiro A.S., Teixeira D.C., Silva A.M., Sardinha L.B., et al. Effects of Whey Protein Supplementation Pre- or Post-Resistance Training on Muscle Mass, Muscular Strength, and Functional Capacity in Pre-Conditioned Older Women: A Randomized Clinical Trial. *Nutrients*. 2018;10:563. doi: 10.3390/nu10050563. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
148. Nabuco H.C.G., Tomeleri C.M., Sugihara Junior P, Fernandes R.R., Cavalcante E.F., Venturini D., Barbosa D.S., Silva A.M., Sardinha L.B., Cyrino E.S. Effects of pre- or post-exercise whey protein supplementation on body fat and metabolic and inflammatory profile in pre-conditioned older women: A randomized, double-blind, placebo-controlled trial. *Nutr. Metab. Cardiovasc. Dis. NMCD*. 2019;29:290–300. doi: 10.1016/j.numecd.2018.11.007. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
149. Nilsson M.I., Mikhail A., Lan L., Di Carlo A., Hamilton B., Barnard K., Hettinga B.P, Hatcher E., Tarnopolsky M.G., Nederveen J.P, et al. A Five-Ingredient Nutritional Supplement and Home-Based Resistance Exercise Improve Lean Mass and Strength in Free-Living Elderly. *Nutrients*. 2020;12:2391. doi: 10.3390/nu12082391. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
150. Oesen S., Halper B., Hofmann M., Jandrasits W., Franzke B., Strasser E.M., Graf A., Tschan H., Bachl N., Quittan M., et al. Effects of elastic band resistance training and nutritional supplementation on physical performance of institutionalised elderly—A randomized controlled trial. *Exp. Gerontol*. 2015;72:99–108. doi: 10.1016/j.exger.2015.08.013. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
151. Orsatti F.L., Maestá N., de Oliveira E.P, Nahas Neto J., Burini R.C., Nunes P.R.P, Souza A.P, Martins F.M., Nahas E.P Adding Soy Protein to Milk Enhances the Effect of Resistance Training on Muscle Strength in Postmenopausal Women. *J. Diet. Suppl*. 2018;15:140–152. doi: 10.1080/19390211.2017.1330794. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
152. Roschel H., Hayashi A.P, Fernandes A.L., Jambassi-Filho J.C., Hevia-Larraín V, de Capitani M., Santana D.A., Gonçalves L.S., de Sá-Pinto A.L., Lima F.R., et al. Supplement-based nutritional strategies to tackle frailty: A multifactorial, double-blind, randomized placebo-controlled trial. *Clin. Nutr*. 2021;40:4849–4858. doi: 10.1016/j.clnu.2021.06.024. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
153. Seino S., Sumi K., Narita M., Yokoyama Y., Ashida K., Kitamura A., Shinkai S. Effects of Low-Dose Dairy Protein Plus Micronutrient Supplementation during Resistance Exercise on Muscle Mass and Physical Performance in Older Adults: A Randomized, Controlled Trial. *J. Nutr. Health Aging*. 2018;22:59–67. doi: 10.1007/s12603-017-0904-5. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
154. Shahar S., Kamaruddin N.S., Badrasawi M., Sakian N.I., Abd Manaf Z., Yassin Z., Joseph L. Effectiveness of exercise and protein supplementation intervention on body composition, functional fitness, and oxidative stress among elderly Malays with sarcopenia. *Clin. Interv. Aging*. 2013;8:1365–1375. doi: 10.2147/CIA.S46826. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
155. Shenoy S., Bedi R., Sandhu J.S. Effect of soy isolate protein and resistance exercises on muscle performance and bone health of osteopenic/osteoporotic post-menopausal women. *J Women Aging*. 2013;25:183–198. doi: 10.1080/08952841.2013.764252. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

156. Soares A.L.S., Machado-Lima A., Brech G.C., Greve J.M.D., Dos Santos J.R., Inojosa T.R., Rogero M.M., Salles J.E.N., Santarem-Sobrinho J.M., Davis C.L., et al. The Influence of Whey Protein on Muscle Strength, Glycemic Control and Functional Tasks in Older Adults with Type 2 Diabetes Mellitus in a Resistance Exercise Program: Randomized and Triple Blind Clinical Trial. *Int. J. Environ. Res. Public Health*. 2023;20:5891. doi: 10.3390/ijerph20105891. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

157. Sugihara Junior P, Ribeiro A.S., Nabuco H.C.G., Fernandes R.R., Tomeleri C.M., Cunha P.M., Venturini D., Barbosa D.S., Schoenfeld B.J., Cyrino E.S. Effects of Whey Protein Supplementation Associated With Resistance Training on Muscular Strength, Hypertrophy, and Muscle Quality in Preconditioned Older Women. *Int. J. Sport Nutr. Exerc. Metab.* 2018;28:528–535. doi: 10.1123/ijsnem.2017-0253. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

158. Tang W, Yan X., Fan P, Xu D., Li Y. Analysis of the Clinical Effect of Dietary Nutrition Intervention Combined with Resistance Training on the Cognitive Function of Elderly Male Patients with Sarcopenia. *Progr. Mod. Biomed.* 2020;20:2698–2701. doi: 10.13241/j.cnki.pmb.2020.14.021. [[DOI](#)] [[Google Scholar](#)]

159. Thomson R.L., Brinkworth G.D., Noakes M., Buckley J.D. Muscle strength gains during resistance exercise training are attenuated with soy compared with dairy or usual protein intake in older adults: A randomized controlled trial. *Clin. Nutr.* 2016;35:27–33. doi: 10.1016/j.clnu.2015.01.018. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

160. Tieland M., Dirks M.L., van der Zwaluw N., Verdijk L.B., van de Rest O., de Groot L.C., van Loon L.J. Protein supplementation increases muscle mass gain during prolonged resistance-type exercise training in frail elderly people: A randomized, double-blind, placebo-controlled trial. *J. Am. Med. Dir. Assoc.* 2012;13:713–719. doi: 10.1016/j.jamda.2012.05.020. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

161. Trevisan M.C., Souza J.M., Marucci Mde F. Influence of soy protein intake and weight training on the resting energy expenditure of postmenopausal women. *Rev. Assoc. Med. Bras.* 2010;56:572–578. doi: 10.1590/S0104-42302010000500020. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

162. Unterberger S., Aschauer R., Zöhrer P.A., Draxler A., Franzke B., Strasser E.M., Wagner K.H., Wessner B. Effects of an increased habitual dietary protein intake followed by resistance training on fitness, muscle quality and body composition of seniors: A randomised controlled trial. *Clin. Nutr.* 2022;41:1034–1045. doi: 10.1016/j.clnu.2022.02.017. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

163. Verdijk L.B., Jonkers R.A.M., Gleeson B.G., Beelen M., Meijer K., Savelberg H.H.C.M., Wodzig K.W.H.W., Dendale P, Van Loon L.J.C. Protein supplementation before and after exercise does not further augment skeletal muscle hypertrophy after resistance training in elderly men. *Am. J. Clin. Nutr.* 2009;89:608–616. doi: 10.3945/ajcn.2008.26626. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

164. Verreijen A.M., Verlaan S., Engberink M.F, Swinkels S., de Vogel-van den Bosch J., Weijs P.J. A high whey protein-, leucine-, and vitamin D-enriched supplement preserves muscle mass during intentional weight loss in obese older adults: A double-blind randomized controlled trial. *Am. J. Clin. Nutr.* 2015;101:279–286. doi: 10.3945/ajcn.114.090290. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

165. Vijayakumaran R.K., Daly R.M., Tan V.P.S. “We Want More”-Perspectives of Sarcopenic Older Women on the Feasibility of High-Intensity Progressive Resistance Exercises and a Whey-Protein Nutrition Intervention. *Front. Nutr.* 2023;10:1176523. doi: 10.3389/fnut.2023.1176523. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
166. Vikberg S., Sörlén N., Brandén L., Johansson J., Nordström A., Hult A., Nordström P. Effects of Resistance Training on Functional Strength and Muscle Mass in 70-Year-Old Individuals With Pre-sarcopenia: A Randomized Controlled Trial. *J. Am. Med. Dir. Assoc.* 2019;20:28–34. doi: 10.1016/j.jamda.2018.09.011. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
167. Villanueva M.G., He J., Schroeder E.T. Periodized resistance training with and without supplementation improve body composition and performance in older men. *Eur. J. Appl. Physiol.* 2014;114:891–905. doi: 10.1007/s00421-014-2821-1. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
168. Weisgarber K.D., Candow D.G., Farthing J.P. Whey protein and high-volume resistance training in postmenopausal women. *J. Nutr. Health Aging.* 2015;19:511–517. doi: 10.1007/s12603-015-0454-7. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
169. Yamada M., Kimura Y., Ishiyama D., Nishio N., Otobe Y., Tanaka T., Ohji S., Koyama S., Sato A., Suzuki M., et al. Synergistic effect of bodyweight resistance exercise and protein supplementation on skeletal muscle in sarcopenic or dynapenic older adults. *Geriatr. Gerontol. Int.* 2019;19:429–437. doi: 10.1111/ggi.13643. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
170. Zdzieblik D., Jendricke P., Oesser S., Gollhofer A., König D. The Influence of Specific Bioactive Collagen Peptides on Body Composition and Muscle Strength in Middle-Aged, Untrained Men: A Randomized Controlled Trial. *Int. J. Environ. Res. Public Health.* 2021;18:4837. doi: 10.3390/ijerph18094837. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
171. Zhao T., Gao H.L., Guo N.Z., Chea Z.M., Wu X.Y., He D.L., Wei L.Q., Shen X.C., Lv W.Y. Effect of health education and whey protein supplementation combined with resistance exercise on the intervention of middle—Aged and elderly people with sarcopenia. *J. Clin. Exp. Med.* 2022;12:90–94. doi: 10.3969/j.issn.1671-4695.2022.01.025. [[DOI](#)] [[Google Scholar](#)]
172. Zhu L.Y., Chan R., Kwok T., Cheng K.C., Ha A., Woo J. Effects of exercise and nutrition supplementation in community-dwelling older Chinese people with sarcopenia: A randomized controlled trial. *Age Ageing.* 2019;48:220–228. doi: 10.1093/ageing/afy179. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
173. Gorissen S.H.M., Trommelen J., Kouw I.W.K., Holwerda A.M., Pennings B., Groen B.B.L., Wall B.T., Churchward-Venne T.A., Horstman A.M.H., Koopman R., et al. Protein Type, Protein Dose, and Age Modulate Dietary Protein Digestion and Phenylalanine Absorption Kinetics and Plasma Phenylalanine Availability in Humans. *J. Nutr.* 2020;150:2041–2050. doi: 10.1093/jn/nxaa024. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
174. Burd N.A., Gorissen S.H., van Vliet S., Snijders T., van Loon L.J. Differences in postprandial protein handling after beef compared with milk ingestion during postexercise recovery: A randomized

controlled trial. *Am. J. Clin. Nutr.* 2015;102:828–836. doi: 10.3945/ajcn.114.103184. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

175. Tang J.E., Moore D.R., Kujbida G.W., Tarnopolsky M.A., Phillips S.M. Ingestion of whey hydrolysate, casein, or soy protein isolate: Effects on mixed muscle protein synthesis at rest and following resistance exercise in young men. *J. Appl. Physiol.* 2009;107:987–992. doi: 10.1152/jappphysiol.00076.2009. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

176. Wilkinson S.B., Tarnopolsky M.A., Macdonald M.J., Macdonald J.R., Armstrong D., Phillips S.M. Consumption of fluid skim milk promotes greater muscle protein accretion after resistance exercise than does consumption of an isonitrogenous and isoenergetic soy-protein beverage. *Am. J. Clin. Nutr.* 2007;85:1031–1040. doi: 10.1093/ajcn/85.4.1031. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

177. Cheng H., Kong J., Underwood C., Petocz P., Hirani V., Dawson B., O’Leary F. Systematic review and meta-analysis of the effect of protein and amino acid supplements in older adults with acute or chronic conditions. *Br. J. Nutr.* 2018;119:527–542. doi: 10.1017/S0007114517003816. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

178. Liao C.D., Lee P.H., Hsiao D.J., Huang S.W., Tsao J.Y., Chen H.C., Liou T.H. Effects of Protein Supplementation Combined with Exercise Intervention on Frailty Indices, Body Composition, and Physical Function in Frail Older Adults. *Nutrients.* 2018;10:1916. doi: 10.3390/nu10121916. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

179. Luo D., Lin Z., Li S., Liu S.-J. Effect of nutritional supplement combined with exercise intervention on sarcopenia in the elderly: A meta-analysis. *Int. J. Nurs. Sci.* 2017;4:389–401. doi: 10.1016/j.ijnss.2017.09.004. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

180. Naclerio E., Larumbe-Zabala E. Effects of Whey Protein Alone or as Part of a Multi-ingredient Formulation on Strength, Fat-Free Mass, or Lean Body Mass in Resistance-Trained Individuals: A Meta-analysis. *Sports Med.* 2016;46:125–137. doi: 10.1007/s40279-015-0403-y. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

181. Liao C.D., Chen H.C., Huang S.W., Liou T.H. The Role of Muscle Mass Gain Following Protein Supplementation Plus Exercise Therapy in Older Adults with Sarcopenia and Frailty Risks: A Systematic Review and Meta-Regression Analysis of Randomized Trials. *Nutrients.* 2019;11:1713. doi: 10.3390/nu11081713. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

182. Gade J., Pedersen R.J., Beck A.M. Effect of Protein or Essential Amino Acid Supplementation During Prolonged Resistance Exercise Training in Older Adults on Body Composition, Muscle Strength, and Physical Performance Parameters: A Systematic Review. *Rehabil. Process Outcome.* 2018;7:1179572718765760. doi: 10.1177/1179572718765760. [[DOI](#)] [[Google Scholar](#)]

183. Hidayat K., Chen G.-C., Wang Y., Zhang Z., Dai X., Szeto I.M.Y., Qin L.-Q. Effects of milk proteins supplementation in older adults undergoing resistance training: A meta-analysis of randomized control trials. *J. Nutr. Health Aging.* 2018;22:237–245. doi: 10.1007/s12603-017-0899-y. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

184. Kim J., Jeong K., Lim S., Lee S., Baek Y. Association of Dietary Protein Sources and Their Adequacy, Body Composition and Risk of Sarcopenic Obesity in South Korean Populations: A Cross-Sectional Study. *Metabolites*. 2024;14:130. doi: 10.3390/metabo14020130. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
185. Kim J., Baek Y., Jeong K., Lee S. Association of Dietary Factors With Grip Strength, Body Fat, and Prevalence of Sarcopenic Obesity in Rural Korean Elderly With Cardiometabolic Multimorbidity. *Front. Nutr.* 2022;9:910481. doi: 10.3389/fnut.2022.910481. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
186. Li S.Y., Lu Z.H., Leung J.C.S., Kwok T.C.Y. Association of dietary protein intake, inflammation with muscle mass, physical performance and incident sarcopenia in Chinese community-dwelling older adults. *J. Nutr. Health Aging*. 2024;28:100163. doi: 10.1016/j.jnha.2024.100163. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
187. Smeuninx B., McKendry J., Wilson D., Martin U., Breen L. Age-Related Anabolic Resistance of Myofibrillar Protein Synthesis Is Exacerbated in Obese Inactive Individuals. *J. Clin. Endocrinol. Metab.* 2017;102:3535–3545. doi: 10.1210/jc.2017-00869. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
188. Kramer I.F., Verdijk L.B., Hamer H.M., Verlaan S., Luiking Y.C., Kouw I.W.K., Senden J.M., van Kranenburg J., Gijsen A.P., Bierau J., et al. Both basal and post-prandial muscle protein synthesis rates, following the ingestion of a leucine-enriched whey protein supplement, are not impaired in sarcopenic older males. *Clin. Nutr.* 2017;36:1440–1449. doi: 10.1016/j.clnu.2016.09.023. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
189. Nunes E.A., Colenso-Semple L., McKellar S.R., Yau T., Ali M.U., Fitzpatrick-Lewis D., Sherifali D., Gaudichon C., Tomé D., Atherton P.J., et al. Systematic review and meta-analysis of protein intake to support muscle mass and function in healthy adults. *J. Cachexia Sarcopenia Muscle*. 2022;13:795–810. doi: 10.1002/jcsm.12922. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
190. Herda A.A., McKay B.D., Herda T.J., Costa P.B., Stout J.R., Cramer J.T. Changes in Strength, Mobility, and Body Composition Following Self-Selected Exercise in Older Adults. *J. Aging Phys. Act.* 2021;29:17–26. doi: 10.1123/japa.2019-0468. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
191. Mallinson J.E., Wardle S.L., O’Leary T.J., Greeves J.P., Cegielski J., Bass J., Brook M.S., Wilkinson D.J., Smith K., Atherton P.J., et al. Protein dose requirements to maximize skeletal muscle protein synthesis after repeated bouts of resistance exercise in young trained women. *Scand. J. Med. Sci. Sports*. 2023;33:2470–2481. doi: 10.1111/sms.14506. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
192. Churchward-Venne T.A., Pinckaers P.J.M., Smeets J.S.J., Betz M.W., Senden J.M., Goessens J.P.B., Gijsen A.P., Rollo I., Verdijk L.B., van Loon L.J.C. Dose-response effects of dietary protein on muscle protein synthesis during recovery from endurance exercise in young men: A double-blind randomized trial. *Am. J. Clin. Nutr.* 2020;112:303–317. doi: 10.1093/ajcn/nqaa073. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

193. Holwerda A.M., Paulussen K.J.M., Overkamp M., Goessens J.P.B., Kramer I.F., Wodzig W., Verdijk L.B., van Loon L.J.C. Dose-Dependent Increases in Whole-Body Net Protein Balance and Dietary Protein-Derived Amino Acid Incorporation into Myofibrillar Protein During Recovery from Resistance Exercise in Older Men. *J. Nutr.* 2019;149:221–230. doi: 10.1093/jn/nxy263. [[DOI ↗](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar ↗](#)]

194. Mitchell C.J., Milan A.M., Mitchell S.M., Zeng N., Ramzan F., Sharma P., Knowles S.O., Roy N.C., Sjödin A., Wagner K.H., et al. The effects of dietary protein intake on appendicular lean mass and muscle function in elderly men: A 10-wk randomized controlled trial. *Am. J. Clin. Nutr.* 2017;106:1375–1383. doi: 10.3945/ajcn.117.160325. [[DOI ↗](#)] [[PubMed](#)] [[Google Scholar ↗](#)]

Associated Data

This section collects any data citations, data availability statements, or supplementary materials included in this article.

Supplementary Materials

[nutrients-16-00941-s001.zip](#) ^(6MB, zip)

Data Availability Statement

Refer to [Supplementary Materials](#). Raw data available on request.

Articles from Nutrients are provided here courtesy of **Multidisciplinary Digital Publishing Institute (MDPI)**